


Candidate de Sitter Vacua: Construction

based on [ArXiv:2406.13751](https://arxiv.org/abs/2406.13751) with Liam McAllister, Jakob Moritz, and Richard Nally

String Phenomenology 2024 in Padova

June 24, 2024

Andreas Schachner



Collaborators



LIAM MCALLISTER

Cornell University



JAKOB MORITZ

CERN



RICHARD NALLY

Cornell University



A Tale of two Talks



LIAM MCALLISTER

Cornell University

Plan for today:


Construction and validation.

Liam's talk tomorrow:

More details of control analysis.



Open access



Our entire data is publicly available on GitHub!

On top of that, we provide

- **independent python code** to compute e.g. the vacuum energy or corrected volumes
- jupyter notebooks to **validate our solutions** in the approximations explained below
- a **tutorial notebook** to work with the data and to start new calculations by e.g. using **CYTools**
- plotting tools to reproduce some figures from our paper

Everyone can explore our solutions for themselves by using our repository!

https://github.com/AndreasSchachner/kklt_de_sitter_vacua



Upshot

First concrete candidates of de Sitter vacua as envisioned by Kachru, Kallosh, Linde and Trivedi (KKLT) 20 years ago.

IMPORTANT CAVEAT:

These vacua are solutions in a particular leading-order EFT that I will define.
Whether these solutions lift to full string theory remains open.



The setup

Type IIB orientifold compactifications

Setup: Type IIB on CY orientifolds X/\mathcal{F} for a holomorphic and isometric involution $\mathcal{F} : X \rightarrow X$.

Notation: complex structure moduli z^a , $a = 1, \dots, h_-^{2,1}(X)$, Kähler moduli T_A , $A = 1, \dots, h_+^{1,1}(X)$ and axiodilaton τ

We will mainly be interested in the **F-term scalar potential** for these fields

$$V_F = e^K (K^{I\bar{J}} D_I W D_{\bar{J}} \bar{W} - 3 |W|^2) , \quad D_I W = \partial_I W + (\partial_I K) W , \quad K_{I\bar{J}} = \partial_I \partial_{\bar{J}} K$$

The superpotential W is given by [[GVW hep-th/9906070](#), [Witten hep-th/9604030](#)]

$$W(z, \tau, T) = W_{\text{flux}}(z, \tau) + W_{\text{np}}(z, \tau, T) , \quad W_{\text{flux}}(z, \tau) = \sqrt{\frac{2}{\pi}} \int_X (F_3 - \tau H_3) \wedge \Omega(z) , \quad W_{\text{np}}(z, \tau, T) = \sum_D A_D(z, \tau) e^{-\frac{2\pi}{c_D} T_D}$$

The 3-form fluxes have to obey the **D3-tadpole cancellation condition**

$$Q_{\text{flux}} + 2(N_{D3} - N_{\bar{D}3}) = Q_O , \quad Q_{\text{flux}} = \int_X H_3 \wedge F_3$$


where Q_O receives contributions from localised sources like O3/O7-planes or D7-branes.

Recipe for KKLT vacua

The KKLT scenario

[Kachru, Kallosh, Linde, Trivedi [hep-th/0301240](https://arxiv.org/abs/hep-th/0301240)]

The KKLT scenario is a proposal to construct de Sitter vacua in string theory.



CLAIM 1.

Well-controlled SUSY AdS₄ exist in Type IIB flux compactifications with

1. $\langle W_{flux} \rangle \ll 1$, and
2. non-perturbative D-brane instantons.

Explicit examples in
[Demirtas et al. [2107.09064](https://arxiv.org/abs/2107.09064)]

CLAIM 2.

For such a SUSY AdS₄, provided one finds

3. warped deformed conifold [Klebanov, Strassler [hep-th/0007191](https://arxiv.org/abs/hep-th/0007191)]
4. containing some anti-D3 branes [Kachru, Pearson, Verlinde [hep-th/0112197](https://arxiv.org/abs/hep-th/0112197)]
5. in a suitable parameter regime

there are metastable dS₄ vacua.

We provide the first examples fulfilling
conditions 1., 2., 3., 4., and 5.

See also e.g.:

[Moritz et al. [1809.06618](#)]

[Bena et al. [1809.06861](#)]

[Carta et al. [1902.01412](#)]


[Dudas, S. Lüst [1912.09948](#)]

[S. Lüst, Randall [2206.04708](#)]

Recipe for KKLT vacua

Uplift to de Sitter vacua

[Kachru, Kallosh, Linde, Trivedi [hep-th/0301240](#)]



Klebanov-Strassler throats arise in CY compactifications through conifold singularities threaded by 3-form fluxes

$$e^{4A_{IR}} \approx e^{-8\pi K/3n_{cf}g_s M} \sim z_{cf}^{\frac{4}{3}}$$

where M, K are the fluxes threading the S^3 of the deformed conifold.

Control over the α' expansion at the tip of the throat, i.e., small curvature at the bottom of the throat requires $g_s M \gtrsim 1$.

[Klebanov, Strassler [hep-th/0007191](#)]

[Giddings, Kachru, Polchinski [hep-th/0105097](#)]


To achieve **CLAIM 2**, an anti-D3 brane at the tip of the throat provides a positive source of energy which potentially uplifts the AdS minimum to a dS minimum provided

$$V_{\text{KPV}}^{\overline{D3}} = \frac{c}{\mathcal{V}_E^{4/3}}, \quad \Xi = \frac{V_{\text{KPV}}^{\overline{D3}}}{V_F} = \frac{\zeta e^{K_{cs}/3}}{(g_s M)^2} \mathcal{V}_E^{2/3} \frac{z_{cf}^{4/3}}{W_0^2} \sim 1, \quad \zeta \approx 114.037$$

We call vacua satisfying $\Xi \sim 1$ **well-aligned** which are the main targets of this talk!

The anti-D3-brane state at the bottom of the Klebanov-Strassler throat is metastable provided $M > 12$.

[Kachru, Pearson, Verlinde [hep-th/0112197](#)]



Recipe for KKLT vacua

Checklist for KKLT vacua



The point of this talk is to show you how to actually accomplish all this in explicit setups!


Constructing the leading order EFT

The working plan





Constructing the leading order EFT

Scan for Geometries and Orientifolds



We will work with mirror pairs of CY_3 hypersurfaces X, \widetilde{X}
 in toric varieties V, \widetilde{V}
 obtained from triangulations of 4D polytopes Δ°, Δ



Demirtas, Rios-Tascon,
 McAllister [2211.03823](https://arxiv.org/abs/2211.03823)

473,800,776 reflexive polytopes in 4D Kreuzer,
 Skarke (KS) [\[hep-th/0002240\]](https://arxiv.org/abs/hep-th/0002240)

We restrict to \mathbb{Z}_2 -involutions $x \rightarrow -x$ with O3/O7-planes for **trilayer** polytopes
 such that $h_-^{1,1} = h_+^{1,2} = 0$ [\[Moritz 2305.06363\]](https://arxiv.org/abs/2305.06363).

We cancel the D7-tadpole locally giving rise to $\mathfrak{so}(8)$ $\mathcal{N} = 1$ super Yang-Mills
 theory hosted on four-cycles with O7-planes.

In these setups, the D3-tadpole is $Q_0 = h^{1,1} + h^{2,1} + 2$.



Constructing the leading order EFT

The Kähler moduli sector

From previous slides, we recall

$$V = V_F + V_{\text{up}} \ , \quad V_F = e^K (K^{I\bar{J}} D_I W D_{\bar{J}} \bar{W} - 3 W^2) \ , \quad V_{\text{up}} = V_{\text{KPV}}^{\overline{\text{D3}}} \ , \quad W(z, \tau, T) = W_{\text{flux}}(z, \tau) + W_{\text{np}}(z, \tau, T)$$

We work in the leading-order EFT where the Kähler potential and Kähler coordinates are given by

$$K_{1.o.} \approx K_{\text{tree}} + K_{(\alpha')^3} + K_{\text{WSI}} \ , \quad T_A^{\text{l.o.}} \approx T_A^{\text{tree}} + \delta T_A^{(\alpha')^2} + \delta T_A^{\text{WSI}}$$

Here the tree level α' and worldsheet instanton (WSI) corrections amount to

$$K_{1.o.} = -2 \log \left[\frac{1}{6} \kappa_{ABC} t^A t^B t^C - \frac{\zeta(3) \chi(X)}{4(2\pi)^3} + \frac{1}{2(2\pi)^3} \sum_{\mathbf{q} \in \mathcal{M}(X)} \mathcal{N}_{\mathbf{q}} \left(\text{Li}_3 \left((-1)^{\mathbf{r} \cdot \mathbf{q}} e^{-2\pi \mathbf{q} \cdot \mathbf{t}} \right) + 2\pi \mathbf{q} \cdot \mathbf{t} \text{Li}_2 \left((-1)^{\mathbf{r} \cdot \mathbf{q}} e^{-2\pi \mathbf{q} \cdot \mathbf{t}} \right) \right) \right] \ ,$$

$$T_A^{\text{l.o.}} = \frac{1}{2} \kappa_{ABC} t^B t^C - \frac{\chi(D_A)}{24} + \frac{1}{(2\pi)^2} \sum_{\mathbf{q} \in \mathcal{M}(X)} q_i \mathcal{N}_{\mathbf{q}} \text{Li}_2 \left((-1)^{\mathbf{r} \cdot \mathbf{q}} e^{-2\pi \mathbf{q} \cdot \mathbf{t}} \right) + i \int_X C_4 \wedge \omega_A \ .$$

Genus-zero Gopakumar-Vafa invariants $\mathcal{N}_{\mathbf{q}}$ [Gopakumar, Vafa [hep-th/9809187](https://arxiv.org/abs/hep-th/9809187)] can be computed using publicly available code: <https://github.com/ariostas/cygv>

For the moment, we ignore

- string loop corrections, especially $\mathcal{N} = 1$ corrections
- α' corrections to the KPV potential for the anti-D3 brane as derived in [Junghans [2201.03572](https://arxiv.org/abs/2201.03572)] [Hebecker, 3xSchreyer, 2xVenken [2208.02826](https://arxiv.org/abs/2208.02826), [2212.07437](https://arxiv.org/abs/2212.07437), [2402.13311](https://arxiv.org/abs/2402.13311)]
- ... see talk by Liam

See in particular:

[Becker et al. [hep-th/0204254](https://arxiv.org/abs/hep-th/0204254)]

[Robles-Llana et al. [hep-th/0612027](https://arxiv.org/abs/hep-th/0612027), [0707.0838](https://arxiv.org/abs/0707.0838)]

[Cecotti et al. [Int.J.Mod.Phys.A 4 \(1989\) 2475](https://arxiv.org/abs/Int.J.Mod.Phys.A.4.1989.2475)]

[Grimm [0705.3253](https://arxiv.org/abs/0705.3253)]

Constructing the leading order EFT

The flux superpotential

The flux superpotential is given in terms of the **period vector** $\vec{\Pi}$ and the **pre-potential** $F = F(z)$ as

$$W_{\text{flux}}(\tau, z^a) = \sqrt{\frac{2}{\pi}} \int_X (F_3 - \tau H_3) \wedge \Omega(z) = \sqrt{\frac{2}{\pi}} \vec{\Pi}^\top \cdot \Sigma \cdot (\vec{f} - \tau \vec{h}) \quad , \quad \vec{\Pi} = (2F - z^a F_a, F_a, 1, z^a) \quad , \quad F_a = \partial_a F$$

We compute $F(z)$ explicitly at **Large Complex Structure (LCS)** using mirror symmetry following [\[Hosono et al. hep-th/9406055\]](#)

$$F_{\text{poly}}(z) = -\frac{1}{3!} \tilde{\kappa}_{abc} z^a z^b z^c + \frac{1}{2} \tilde{a}_{ab} z^a z^b + \frac{1}{24} \tilde{c}_a z^a + \frac{\zeta(3) \chi(\tilde{X})}{2(2\pi i)^3} \quad , \quad F_{\text{inst}}(z) = -\frac{1}{(2\pi i)^3} \sum_{\tilde{\mathbf{q}} \in \mathcal{M}(\tilde{X})} \mathcal{N}_{\tilde{\mathbf{q}}} \text{Li}_3\left(e^{2\pi i \tilde{\mathbf{q}} \cdot \mathbf{z}}\right)$$

in terms of quantities $\tilde{\kappa}_{abc}, \tilde{a}_{ab}, \tilde{c}_a$ defined on the mirror CY \tilde{X} , see e.g. [\[Demirtas et al. 2303.00757\]](#).

It is known how to construct **conifolds** by shrinking a set of curves in \tilde{X} to zero volume [\[Demirtas et al. 2009.03312\]](#) [\[Álvarez-García et al. 2009.03325\]](#).

We write $z^a = (z_{\text{cf}}, z^\alpha)$, $\alpha = h^{2,1}(X) - 1$, and expand the periods order by order in the conifold modulus z_{cf}

$$W_{\text{flux}}(z^a, \tau) = W_{\text{poly}}(z^\alpha, \tau) + W_{\text{inst}}(z^\alpha, \tau) + z_{\text{cf}} W^{(1)}(z^\alpha, z_{\text{cf}}, \tau) + \mathcal{O}(z_{\text{cf}}^2).$$

Constructing the leading order EFT

The non-perturbative superpotential

The non-perturbative superpotential from D-branes wrapping rigid divisors D reads [Witten [hep-th/9610234](#)]

$$W_{\text{np}}(z, \tau, T) = \sum_D A_D(z, \tau) e^{-\frac{2\pi}{c_D} T_D}, \quad c_D = \begin{cases} 1 & \text{Euclidean D3-branes,} \\ 6 & \text{gaugino condensation on 7-branes.} \end{cases}$$

We check that the **only** contributing divisors are **pure rigid** implying [Witten [hep-th/9610234](#), Demirtas et al. [2107.09064](#)]

$$A_D(z, \tau) = A_D = \text{const}$$

For the normalisation of the A_D we choose

$$A_D = \sqrt{\frac{2}{\pi}} \frac{n_D}{(4\pi)^2}.$$

The constant n_D is

- related to an integral over worldsheet modes [Alexandrov et al. [2204.02981](#)], and
- expected to be an order-one number due to mirror symmetry.

Computing n_D has so far been out of reach.

In our vacua, we take $n_D = 1$ and then check a posteriori that our vacua persist for $10^{-3} \leq n_D \leq 10^4$.

See also

[Kim [2107.09779](#), [2301.03602](#)]

[Jefferson, Kim [2211.00210](#)]

Finding KKLT vacua in KS

The working plan



Finding KKLT vacua in KS

Perturbatively Flat Vacua (PFVs)

[Demirtas, Kim, McAllister, Moritz: [1912.10047](#)]

In the presence of conifolds

[Demirtas et al. [2009.03312](#)]

[Álvarez-García et al. [2009.03325](#)]

For special flux choices $\vec{M}, \vec{K} \in \mathbb{Z}^{h^{2,1}}$, the polynomial flux superpotential W_{poly} and the F-terms vanish along $z^a = p^a \tau$ where

$$p^a = (N^{-1})^{ab} K_b, \quad N_{ab} = \tilde{\kappa}_{abc} M^c$$

The remaining superpotential terms are computable in terms of GV invariants on \tilde{X}

$$W_{\text{inst}} = \frac{-1}{(2\pi)^2} \sum_{\tilde{\mathbf{q}} \in \mathcal{M}(\tilde{X})} \mathcal{N}_{\tilde{\mathbf{q}}} (M^a \tilde{\mathbf{q}}_a) \text{Li}_2(e^{2\pi i \tilde{\mathbf{q}}_a p^a \tau})$$

A minimum for the light degree of freedom τ arises frequently through the **racetrack mechanism** so that

$$W_0 = \langle W_{\text{flux}} \rangle = \langle W_{\text{inst}} \rangle \ll 1$$

In practice, we obtain the **true minimum** by numerically solving F-term conditions.

For related work, see also

[Honma, Otsuka [2103.03003](#)]

[Marchesano et al. [2105.09326](#)]

[Broeckel et al. [2108.04266](#)]


[Basitjan et al. [2108.11962](#)]

[Carta et al. [2112.13863](#)]

[Blumenhagen et al. [2206.08400](#)]

[Cicoli et al. [2209.02720](#)]

Finding KKLT vacua in KS



Kähler moduli stabilisation in explicit setups

Demirtas, Kim, McAllister, Moritz, Rios-Tascon: [2107.09064](#)

To solve the F-terms for the Kähler moduli,

$$D_A W = \partial_A W + K_A W = 0,$$

we use an algorithm described in [\[Demirtas et al. 2107.09064\]](#)

1. Pick arbitrary triangulation of Δ° and choose arbitrary point t_0^A in the Kähler cone
2. Find **initial guess** as classical F-term minimum


$$T_A^0 = \frac{1}{2} \kappa_{ABC} t_0^B t_0^C \rightarrow T_A^* \approx \frac{c_A}{2\pi} \log(W_0^{-1})$$

2. Obtain **true F-term minimum including corrections** by using e.g. Newton's method


$$T_A^* \rightarrow \langle T_A \rangle$$

In the absence of conifolds, this was achieved explicitly in [\[Demirtas et al. 2107.09064, 2107.09065\]](#).

We have new solutions with KS throats and only even fluxes [\[McAllister, Moritz, Nally, AS: 2406.13751\]](#).




Extended Kähler cone



Finding KKLT vacua in KS

De Sitter vacuum containing anti-D3 branes

McAllister, Moritz, Nally, AS: [2406.13751](#)



We restrict to configurations with $Q_{\text{flux}} = Q_0 + 2$ for which the tadpole is cancelled exactly by adding a **single anti-D3 brane** at the tip of the throat.

This makes the previous AdS geometry an **unphysical AdS precursor!**

Practically, it is however important because it makes it easier to locate the true uplifted minimum!


The vacuum is obtained by solving

$$\partial_A V = \partial_A (V_F + V_{\text{up}}) = 0, \quad V_{\text{up}} \sim \frac{e^{-8\pi K/3n_{cf}g_s M}}{\mathcal{V}^{4/3}}$$

for the Kähler moduli **and** complex structure moduli. We follow the same strategy as before:

1. Use triangulation and Kähler parameters for the AdS precursor as initial guess
2. Obtain **uplifted non-SUSY (A)dS vacuum** (if it exists) by using Newton's method

$$\langle T_A \rangle_{\text{AdS}} \rightarrow \langle T_A \rangle$$



Extended Kähler cone

Explicit examples of KKLT vacua

The working plan



Let us put everything together ...



Explicit examples of KKLT vacua

The scan for suitable candidates

McAllister, Moritz, Nally, AS: [2406.13751](#)

| Condition | Number of configurations |
|---|----------------------------|
| $3 \leq h^{2,1} \leq 8$ | 202,073 polytopes |
| trilayer, Δ and Δ° favorable | 3187 polytopes |
| Hodge number cuts | 322 polytopes |
| $\geq h^{1,1}$ rigid divisors | 322 polytopes |
| conifold disjoint from O-planes | 2669 conifolds |
| conifold consistent with KKLT point | 416 conifolds |
| fluxes giving conifold PFV | 240,480,253 conifold PFVs |
| two-term racetrack | 141,594,222 racetrack PFVs |
| $M > 12$; one anti-D3-brane | 33,371 anti-D3-brane PFVs |



Explicit examples of KKLT vacua

Racetrack minima with anti-D3 branes

McAllister, Moritz, Nally, AS: [2406.13751](#)

We obtained **33,371** anti-D3 PFVs with $Q_{\text{flux}} = Q_0 + 2$ of which 396 satisfy

$$0.1 \leq \mathbb{E} \leq 10, \quad g_s < 0.4, \quad W_0 < 0.1.$$



Explicit candidates of KKLT vacua

One de Sitter to rule them all

McAllister, Moritz, Nally, AS: [2406.13751](#)


Here is an explicit example of a de Sitter candidate vacuum at $h^{1,1} = 150$ and $h^{2,1} = 8$

$$\vec{M} = (16, 10, -26, 8, 32, 30, 18, 28)^\top, \quad \vec{K} = (-6, -1, 0, 1, -3, 2, 0, -1)^\top, \quad \vec{p} = \frac{1}{40}(0, -8, 0, -2, 4, 5, 5, 4)^\top$$


giving rise to

$$g_s = 0.0657 \quad W_0 = 0.0115 \quad g_s M = 1.051 \quad z_{\text{cf}} = 2.882 \times 10^{-8} \quad V_{\text{dS}} = +1.937 \times 10^{-19} M_{\text{pl}}^4$$

Potential before and after uplift:



The vacuum is free of tachyons!




Explicit candidates of KKLT vacua

One de Sitter to rule them all


McAllister, Moritz, Nally, AS: [2406.13751](#)

$$W_N \sim \sum_{\tilde{\mathbf{q}} \cdot p = N} \mathcal{N}_{\tilde{\mathbf{q}}} (M^a \tilde{\mathbf{q}}_a) \text{Li}_2(e^{2\pi i N \tau})$$



Racetrack potential

$$\xi_n = \mathcal{N}_{n\mathbf{q}} e^{-2\pi n \mathbf{q} \cdot \mathbf{t}}$$



Contributions from potent curves

Pfaffian prefactors:

Recall the for our results, we set $n_D = 1$ in

$$A_D = \sqrt{\frac{2}{\pi}} \frac{1}{(4\pi)^2} \times n_D$$

We checked that our vacua survive for


$$10^{-3} \leq n_D \leq 10^4$$

See Liam's talk...





Explicit candidates of KKLT vacua

One **five de Sitters** to rule them all

McAllister, Moritz, Nally, AS: [2406.13751](#)



| ID | $h^{2,1}$ | $h^{1,1}$ | M | K' | g_s | W_0 | $g_s M$ | $ z_{cf} $ | V_0 |
|----|-----------|-----------|-----|-----------------|--------|---------|---------|------------------------|--------------------------|
| 1 | 8 | 150 | 16 | $\frac{26}{5}$ | 0.0657 | 0.0115 | 1.051 | 2.822×10^{-8} | $+1.937 \times 10^{-19}$ |
| 2 | 8 | 150 | 16 | $\frac{93}{19}$ | 0.0571 | 0.00490 | 0.913 | 7.934×10^{-9} | $+1.692 \times 10^{-20}$ |
| 3 | 8 | 150 | 18 | $\frac{40}{11}$ | 0.0442 | 0.0222 | 0.796 | 8.730×10^{-8} | $+4.983 \times 10^{-19}$ |
| 4 | 5 | 93 | 20 | $\frac{17}{5}$ | 0.0404 | 0.0539 | 0.808 | 1.965×10^{-6} | $+2.341 \times 10^{-15}$ |
| 5 | 5 | 93 | 16 | $\frac{29}{10}$ | 0.0466 | 0.0304 | 0.746 | 8.703×10^{-7} | $+2.113 \times 10^{-15}$ |



Conclusions

Main takeaway:

First explicit candidate de Sitter solutions along the lines anticipated by Kachru, Kallosh, Linde and Trivedi in '03.

The control parameters in our solutions are currently the best we could do in 2024, but we barely scratched the surface of available compactifications in the KS database.

Open issues and future directions:

- dS vacua are probably most vulnerable to corrections to the anti-D3 brane,
- meta-stability of the uplift in the regime $g_s M \sim 1$ remains an important open problem!
- better understanding the structure of corrections (like string loop or warping corrections),
- perturbations of the throat (would require computing the CY-metric), and
- flux quantisation conditions for CY orientifolds (for toroidal orientifolds, some fluxes have to be even)

[Junghans [2201.03572](#)]

[Hebecker, Schreyer, Venken [2208.02826](#)]

[Schreyer, Venken [2212.07437](#)]

[Schreyer [2402.13311](#)]

[Frey, Polchinski [hep-th/0201029](#)]

In the future, some candidate vacua may survive as genuine de Sitter vacua of string theory.



A wide-angle landscape photograph featuring a large, calm lake in the foreground. The lake's surface is dark blue-green and reflects the bright sun, which is positioned high in the sky, creating a prominent lens flare and a shimmering path of light on the water. The middle ground is dominated by a thick layer of white mist or fog that fills the valley, partially obscuring a small town with white buildings and dark roofs on the left. In the background, several mountain ranges are visible, with some peaks covered in snow and others in dark green forest. The sky is a clear, vibrant blue with a few wispy white clouds. The overall atmosphere is serene and peaceful.


Thank you!



Finding KKLT vacua in KS

A landscape of supersymmetric AdS vacua

McAllister, Moritz, Nally, AS: [2406.13751](https://arxiv.org/abs/2406.13751), work in progress




We found new supersymmetric AdS vacua with **Klebanov-Strassler throats**...

| ID | $(h^{2,1}, h^{1,1})$ | M | K' | N_{D3} | g_s | W_0 | $g_s M$ | $ z_{cf} $ | $-V_F$ | Ξ |
|----|----------------------|-----|----------------|----------|-------------------|----------------------|---------|----------------------|----------------------|-----------|
| a | (6, 160) | 8 | $\frac{1}{15}$ | 2 | $3 \cdot 10^{-3}$ | $1.0 \cdot 10^{-35}$ | 0.021 | $6.0 \cdot 10^{-6}$ | $2.5 \cdot 10^{-90}$ | 10^{70} |
| b | (7, 155) | 8 | 2 | 0 | 0.18 | $7.4 \cdot 10^{-18}$ | 1.46 | $2.1 \cdot 10^{-3}$ | $5.1 \cdot 10^{-50}$ | 10^{34} |
| c | (6, 160) | 2 | 10 | 0 | 0.015 | $1.6 \cdot 10^{-27}$ | 0.30 | $2.4 \cdot 10^{-47}$ | $5.8 \cdot 10^{-72}$ | 0.06 |
| d | (6, 160) | 2 | $\frac{33}{2}$ | 11 | 0.27 | $3.2 \cdot 10^{-25}$ | 0.55 | $1.3 \cdot 10^{-42}$ | $2.3 \cdot 10^{-66}$ | 0.65 |
| e | (8, 150) | 14 | 4 | 0 | 0.075 | 0.032 | 1.05 | $9.1 \cdot 10^{-7}$ | $1.8 \cdot 10^{-17}$ | 3.38 |

Can add brane-antibrane pair and achieve uplift to positive energy.

Interesting candidate for an explicit setting for the inflationary scenario of [KKLMMT: [hep-th/0308055](https://arxiv.org/abs/hep-th/0308055)]

Finding KKLT vacua in KS



A landscape of supersymmetric AdS vacua

McAllister, Moritz, Nally, AS: [2406.13751](https://arxiv.org/abs/2406.13751), work in progress


We found new supersymmetric AdS vacua with **Klebanov-Strassler throats**...

| ID | $(h^{2,1}, h^{1,1})$ | M | K' | N_{D3} | g_s | W_0 | $g_s M$ | $ z_{cf} $ | $-V_F$ | Ξ |
|----|----------------------|-----|----------------|----------|-------------------|----------------------|---------|----------------------|----------------------|-----------|
| a | (6, 160) | 8 | $\frac{1}{15}$ | 2 | $3 \cdot 10^{-3}$ | $1.0 \cdot 10^{-35}$ | 0.021 | $6.0 \cdot 10^{-6}$ | $2.5 \cdot 10^{-90}$ | 10^{70} |
| b | (7, 155) | 8 | 2 | 0 | 0.18 | $7.4 \cdot 10^{-18}$ | 1.46 | $2.1 \cdot 10^{-3}$ | $5.1 \cdot 10^{-50}$ | 10^{34} |
| c | (6, 160) | 2 | 10 | 0 | 0.015 | $1.6 \cdot 10^{-27}$ | 0.30 | $2.4 \cdot 10^{-47}$ | $5.8 \cdot 10^{-72}$ | 0.06 |
| d | (6, 160) | 2 | $\frac{33}{2}$ | 11 | 0.27 | $3.2 \cdot 10^{-25}$ | 0.55 | $1.3 \cdot 10^{-42}$ | $2.3 \cdot 10^{-66}$ | 0.65 |
| e | (8, 150) | 14 | 4 | 0 | 0.075 | 0.032 | 1.05 | $9.1 \cdot 10^{-7}$ | $1.8 \cdot 10^{-17}$ | 3.38 |

... and in addition with **only even fluxes**.


| ID | $(h^{2,1}, h^{1,1})$ | M | K' | N_{D3} | g_s | W_0 | $g_s M$ | $ z_{cf} $ | V_F | Ξ |
|----|----------------------|-----|----------------|----------|-------|----------------------|---------|----------------------|-----------------------|-----------|
| f | (7, 155) | 8 | 2 | 0 | 0.18 | $9.7 \cdot 10^{-18}$ | 1.46 | $2.1 \cdot 10^{-3}$ | $-8.3 \cdot 10^{-50}$ | 10^{35} |
| g | (8, 150) | 8 | $\frac{54}{7}$ | 0 | 0.23 | $2.3 \cdot 10^{-2}$ | 1.86 | $3.1 \cdot 10^{-7}$ | $-1.6 \cdot 10^{-16}$ | 0.28 |
| h | (6, 160) | 4 | $\frac{7}{2}$ | -2 | 0.056 | $1.9 \cdot 10^{-11}$ | 0.23 | $1.0 \cdot 10^{-22}$ | $-2.2 \cdot 10^{-38}$ | 0.22 |

Requires two anti-D3 branes



Racetrack PFVs

McAllister, Moritz, Nally, AS: [2406.13751](#)



We see that both $M \gg 1$ and KS throats containing almost the entire D3-brane charge of the compactification occur in our ensemble, but **both are exponentially rare.**

