


(2)

(1)



# A new modular mechanism for neutrino masses from low-scale seesaw

Matteo Parriciatu <sup>(1)(2)</sup>

from a work in preparation with A. Granelli, D. Meloni, J.T. Penedo, S.T. Petcov



# PLANCK 2025

May 27th, 2025 Padova



## 3 $\nu$ Paradigm

$$\underbrace{|\nu_\alpha\rangle}_{\text{Interaction basis}} = \sum_{i=1}^3 U_{\alpha i}^* \underbrace{|\nu_i\rangle}_{\text{Mass basis}}$$

$$\begin{array}{ccc} \theta_{12} & \theta_{13} & \theta_{23} \\ \delta_{CP} & \boxed{\alpha_{21}, \alpha_{31}} & \end{array}$$

Majorana

$$U_{\text{PMNS}} = \begin{pmatrix} c_{12}c_{13} & s_{12}c_{13} & s_{13}e^{-i\delta_{CP}} \\ -s_{12}c_{23} - c_{12}s_{13}s_{23}e^{i\delta_{CP}} & c_{12}c_{23} - s_{12}s_{13}s_{23}e^{i\delta_{CP}} & c_{13}s_{23} \\ s_{12}s_{23} - c_{12}s_{13}c_{23}e^{i\delta_{CP}} & -c_{12}s_{23} - s_{12}s_{13}c_{23}e^{i\delta_{CP}} & c_{13}c_{23} \end{pmatrix} \times \text{diag}(1, e^{i\alpha_{21}}, e^{i\alpha_{31}})$$

$$c_{ij} \equiv \cos \theta_{ij}, \quad s_{ij} \equiv \sin \theta_{ij} \quad \text{Pontecorvo - Maki - Nakagawa - Sakata}$$

$$P(\nu_\alpha \rightarrow \nu_\beta) = \sum_{i=1}^3 \sum_{j=1}^3 U_{\alpha i}^* U_{\beta i} U_{\alpha j} U_{\beta j}^* \exp \left( -i \frac{\Delta m_{ij}^2 L}{2E} \right)$$

Sign determined from MSW


$$r \equiv \frac{\Delta m_{\text{sol}}^2}{|\Delta m_{\text{atm}}^2|} \approx 1/30$$

Undetermined sign

$$|\Delta m_{\text{sol}}^2| \equiv |m_2^2 - m_1^2|$$

$$|\Delta m_{\text{atm}}^2| \equiv |m_3^2 - m_{1,2}^2|$$

?



## Best fit values



$$\theta_{12} \approx 33.4^\circ, \quad \theta_{13} \approx 8.5^\circ, \quad \theta_{23} \approx 42.4^\circ \quad (\text{NO})$$

$$\theta_{12} \approx 33.4^\circ, \quad \theta_{13} \approx 8.5^\circ, \quad \theta_{23} \approx 48.9^\circ \quad (\text{IO})$$

Two big and  
one small  
angles

F. Capozzi et al.

*Phys. Rev. D* **104** (Oct, 2021)



$3\sigma$  ranges from

F. Capozzi et al.

*Phys. Rev. D* **104** (Oct, 2021)


$$J_{CP} = c_{12}s_{12}c_{23}s_{23}c_{13}^2s_{13}\sin\delta_{CP}$$

Violation if

$$\frac{\delta_{CP}}{\pi} \neq \{0, 1\}$$

CP-conserving still allowed  
at  $2\sigma$  (NO)





$$\sum_i^3 m_i < 0.115 \text{ eV} \quad (95\% \text{ C.L.})$$

A. Shadab et al.

*Phys. Rev. D* **103** (Apr, 2021)


Cosmology

$$\sqrt{\sum_i m_i^2 |U_{ei}|^2} < 0.45 \text{ eV}$$

*Science* 388 (2025) 6743

KATRIN





$$m_e/m_\mu \simeq \frac{1}{200}$$

$$m_\mu/m_\tau \simeq \frac{1}{17}$$


$$\|V_{CKM}\| = \begin{pmatrix} \text{black} & \text{gray} & \text{white} \\ \text{gray} & \text{black} & \text{white} \\ \text{white} & \text{white} & \text{black} \end{pmatrix} \quad \|U_{PMNS}\| = \begin{pmatrix} \text{black} & \text{gray} & \text{gray} \\ \text{gray} & \text{gray} & \text{gray} \\ \text{gray} & \text{gray} & \text{gray} \end{pmatrix}$$

0 0.2 0.4 0.6 0.8 1

D. Meloni [1709.02662v2]



F. Feruglio  
[1706.08749]



$\tau \equiv \text{modulus}$

$$(y_{jk} \bar{L}_j H \ell_{Rk} + \text{h.c.})$$



Free structureless parameters  
in the SM



**Modular Invariance**



$Y(\tau)$

Predictivity




User-friendly



Modular forms:  
less free parameters





## A substantial amount of successful modular models

For reviews, see

T. Kobayashi, M. Tanimoto (2307.03384)

G.-J. Ding, S.F. King (2311.09282)

Attempts include:

Both lepton and quark sectors



Strong CP problem



(2305.08908)

Inflation






(2405.06497)

Baryogenesis





(2504.03506)





$$\tau \equiv \frac{\omega_2}{\omega_1}$$

Modulus






Change basis?

$$\begin{pmatrix} \omega'_2 \\ \omega'_1 \end{pmatrix} = \gamma \begin{pmatrix} \omega_2 \\ \omega_1 \end{pmatrix} \equiv \begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} \omega_2 \\ \omega_1 \end{pmatrix} \quad a, b, c, d \in \mathbb{Z}$$




$$\gamma \in \mathrm{SL}(2, \mathbb{Z}) \equiv \text{Modular group} \equiv \Gamma$$

$$\tau \equiv \frac{\omega_2}{\omega_1} \xrightarrow{\mathrm{SL}(2, \mathbb{Z})} \tau' = \frac{a \omega_2 + b \omega_1}{c \omega_2 + d \omega_1} = \frac{a\tau + b}{c\tau + d}$$







$$\int d^4x d^6y \mathcal{L}_{10D} \implies \int d^4x \mathcal{L}_{EFT}$$

What is a modular form?  $Y(\tau)$

►  $Y(\gamma(\tau)) = (c\tau + d)^k Y(\tau)$

Holomorphic in:  $\{\tau \in \mathbb{C} \mid \text{Im}(\tau) > 0\}$


► “Weight”  $k > 0$

$$c, d \in \begin{pmatrix} a & b \\ c & d \end{pmatrix} \equiv \gamma$$

$$a, b, c, d \in \mathbb{Z} \quad , \quad ad - bc = 1$$

Very  
constraining!





## Superfields transformations

$$\begin{cases} \tau \rightarrow \gamma(\tau) = \frac{a\tau + b}{c\tau + d} & \gamma \in \Gamma \\ \varphi^{(I)} \rightarrow (c\tau + d)^{-k_I} \underbrace{\rho^{(I)}(\gamma)} \varphi^{(I)} \end{cases}$$

Usual matter fields

Unitary irrep. of  $\Gamma_N \subset \Gamma$

$N = 1, 2, 3, \dots$  "Level"

## Finite modular group

$$\Gamma_N$$

for  $N \leq 5$  isomorphic to

$$N = 2$$

$$S_3$$

$$N = 3$$

$$A_4$$


$$N = 4$$

$$S_4$$

$$N = 5$$

$$A_5$$

non-abelian discrete groups





►  $\mathcal{W}(\Phi) = \sum (Y_{I_1 \dots I_n}(\tau) \varphi^{(I_1)} \dots \varphi^{(I_n)})_1$        $\mathcal{W}(\Phi)$  modular invariant if:

$$\begin{cases} \rho \otimes \rho_{I_1} \otimes \rho_{I_2} \dots \otimes \rho_{I_n} \supset \mathbf{1} \\ k_Y = k_{I_1} + k_{I_2} + \dots + k_{I_n} \end{cases} \begin{matrix} \longrightarrow \\ \longrightarrow \end{matrix} \begin{matrix} \text{Usual} \\ \text{Novelty} \end{matrix}$$

$$Y_{I_1 \dots I_n}(\tau) \rightarrow (c\tau + d)^{k_Y} \rho(\gamma) Y_{I_1 \dots I_n}(\tau)$$


Yukawa: modular forms of weight  $k_Y$

$$\varphi^{(I)} \rightarrow (c\tau + d)^{-k_I} \rho^{(I)}(\gamma) \varphi^{(I)}$$

Superfields with modular charges  $-k_I$

$$(c\tau + d)^{k_Y} (c\tau + d)^{-\sum k_{I_n}} = 1$$

## Symmetry breaking



$$Y(\tau)$$

uniquely determined by  $\tau$

$$\text{Fourier} \sim \sum_n^{\infty} a_n q^n$$


$$q \equiv e^{2\pi i \tau}$$



## Lepton mass matrices

$$M_e \sim \sum_i \alpha_i \begin{pmatrix} f_{11}(\tau) & f_{12}(\tau) & \dots \\ \dots & \dots & \dots \\ \dots & \dots & f_{33}(\tau) \end{pmatrix}$$

$f_{ij} \equiv$  pre-determined functions of  $\tau$



$\alpha_i \equiv$  limited number of free parameters ✓



Impose CP symmetry on the model

► gCP  $\Rightarrow \alpha_i \in \mathbb{R}$

P. Novichkov, J. Penedo, S. Petcov, A. Titov


*Journal of High Energy Physics* **2019** no. 7, (Jul, 2019)

Only source of CPV in the model is the VEV of  $\tau$



$$\langle \tau \rangle = \text{Re } \tau + i \text{Im } \tau$$





► Every  $\tau \notin \mathcal{D}$  can be mapped in  $\tau' \in \mathcal{D}$  through  $\Gamma$  transformation


► **CP conserving values**

P. Novichkov, J. Penedo, S. Petcov, A. Titov

*Journal of High Energy Physics* **2019** no. 7, (Jul, 2019)


Only source of CPV in the model is the VEV of  $\tau$

$$\mathcal{D} = \left\{ \tau \in \mathbb{C} : \text{Im } \tau > 0, \quad |\text{Re } \tau| \leq \frac{1}{2}, \quad |\tau| \geq 1 \right\}$$




- Usually the hierarchies put **by hand** (e.g.  $y_e \ll y_\mu \ll y_\tau$ ) in the free parameters
- Froggatt-Nielsen approaches need additional scalars
- **The modular way:** no new scalars, only  $\tau$  needed!

2102.07488 Novichkov, Penedo, Petcov



$$\mathcal{D} = \left\{ \tau \in \mathbb{C} : \text{Im } \tau > 0, \quad |\text{Re } \tau| \leq \frac{1}{2}, \quad |\tau| \geq 1 \right\}$$






► Usually the hierarchies put **by hand** (e.g.  $y_e \ll y_\mu \ll y_\tau$ ) in the free parameters


► Froggatt-Nielsen approaches need **additional scalars**

► **The modular way:** no new scalars, only  $\tau$  needed!

2102.07488 Novichkov, Penedo, Petcov



$$\mathcal{D} = \left\{ \tau \in \mathbb{C} : \text{Im } \tau > 0, \quad |\text{Re } \tau| \leq \frac{1}{2}, \quad |\tau| \geq 1 \right\}$$



- **The modular way:** no new scalars, only  $\tau$  needed!

2102.07488 Novichkov, Penedo, Petcov




- $\tau = \tau_{\text{sym}}$   $\epsilon = |\tau - \tau_{\text{sym}}| > 0$
- $$M \sim \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \rightarrow M \sim \begin{pmatrix} 1 & \epsilon^{\cdots} & \epsilon^{\cdots} \\ \epsilon^{\cdots} & \epsilon^{\cdots} & \epsilon^{\cdots} \\ \epsilon^{\cdots} & \epsilon^{\cdots} & \epsilon^{\cdots} \end{pmatrix}$$

Adapted from J.T.Penedo's slides at DISCRETE 2021

- Mass hierarchies: e.g.  $(1, \epsilon, \epsilon^2)$
- The pattern depends on both  $\Gamma'_N$  and the weights





- ▶ Can “ $\epsilon$ ” also participate in new physics phenomenology?

# Introducing Modular-Symmetry-Protected seesaw



- Can “ $\epsilon$ ” also participate in new physics phenomenology?

## Ingredients:

- The familiar symmetry-protected low-scale type-I seesaw  
see e.g. S. Antusch, O. Fischer 1502.05915

$$\left( \begin{array}{ccc|ccc} 0 & 0 & 0 & y_{11} v_u & 0 & 0 \\ 0 & 0 & 0 & y_{21} v_u & 0 & 0 \\ 0 & 0 & 0 & y_{31} v_u & 0 & 0 \\ \hline y_{11} v_u & y_{21} v_u & y_{31} v_u & 0 & M_N & 0 \\ 0 & 0 & 0 & M_N & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & M_3 \end{array} \right) \quad \text{L-conserving full 6x6 neutrino mass matrix}$$

- $m_\nu = 0$  in this L conserving matrix

## Ingredients:

- The familiar symmetry-protected low-scale type-I seesaw

see e.g. S. Antusch, O. Fischer 1502.05915

$$m_\nu = -m_D M_R^{-1} m_D^T$$

$$\left( \begin{array}{ccc|ccc} 0 & 0 & 0 & y_{11} v_u & 0 & 0 \\ 0 & 0 & 0 & y_{21} v_u & 0 & 0 \\ 0 & 0 & 0 & y_{31} v_u & 0 & 0 \\ \hline y_{11} v_u & y_{21} v_u & y_{31} v_u & 0 & M_N & 0 \\ 0 & 0 & 0 & M_N & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & M_3 \end{array} \right)$$

L-conserving full 6x6  
neutrino mass matrix

- Modular Symmetry mimics L-symmetry in the symmetric points  
 $\tau = \{\omega, i\infty\}$
- L broken by  $\epsilon \equiv |\tau - \tau_{sym}|$



## Ingredients:

- ▶ Modular Symmetry mimics L-symmetry in the symmetric points  $\tau = \{\omega, i\infty\}$
- ▶ Massless neutrinos in the limit of L-symmetry

$$\begin{array}{c} m_0 \\ \left( \begin{array}{ccc|ccc} 0 & 0 & 0 & y_{11} v_u & 0 & 0 \\ 0 & 0 & 0 & y_{21} v_u & 0 & 0 \\ 0 & 0 & 0 & y_{31} v_u & 0 & 0 \\ \hline y_{11} v_u & y_{21} v_u & y_{31} v_u & 0 & M_N & 0 \\ 0 & 0 & 0 & M_N & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & M_3 \end{array} \right) \end{array} \quad \begin{array}{c} M_0 \end{array}$$

▶ pseudo-Dirac pair of mass  $M_N$

## Ingredients:

$$m_\nu = -m_D M_R^{-1} m_D^T$$

- Small neutrino masses controlled also by  $\epsilon \equiv |\tau - \tau_{sym}|$

$$m_D = m_0 + \epsilon^d m_1$$

$$M_R = M_0 + \epsilon^r M_1$$

Controlled by modular forms



## Ingredients:

- Small neutrino masses controlled by  $\epsilon \equiv |\tau - \tau_{sym}|$

$$m_i^\nu \sim \max(|\epsilon|^r, |\epsilon|^d) \times \frac{y^2 v_u^2}{M} \quad y \sim \mathcal{O}(10^{-6}) \text{ and } M \sim \text{GeV}$$

- Mass splitting of the heavy pseudo-Dirac pair also controlled by  $\epsilon \equiv |\tau - \tau_{sym}|$

$$\Delta M \sim |\epsilon|^r \times M. \quad \text{If } M_1 \neq 0$$

$$\Delta M \sim |\epsilon|^d \times \frac{y^2 v_u^2}{M} \quad \text{If } M_1 = 0$$

## The pay-off

- ▶ Charged-leptons mass hierarchies from the same  $\epsilon$  that breaks L-symmetry, gives  $m_\nu \neq 0$  ,  $\Delta M \neq 0$  !
- ▶ Limited number of free parameters
- ▶ HNL phenomenology at colliders (low-scale seesaw) and at intensity frontier



# An example

- ▶ Charged-leptons mass hierarchies controlled by the same  $\epsilon$  which breaks L-symmetry, and gives  $m_\nu \neq 0$  ,  $\Delta M \neq 0$  !

| Model | Group  | $\tau_{\text{sym}}$ | $L$                                                                              | $E^c$              | $N^c$              | $\Delta M$          |
|-------|--------|---------------------|----------------------------------------------------------------------------------|--------------------|--------------------|---------------------|
| A     | $A_4$  | $\omega$            | $(\mathbf{1}, +2) \oplus (\mathbf{1}', +4) \oplus (\mathbf{1}'', +6)$            | $(\mathbf{3}, 0)$  | $(\mathbf{3}, 0)$  | $\sim m_\nu$        |
| B     |        |                     | $(\mathbf{1}', +1) \oplus (\mathbf{1}'', +3) \oplus (\mathbf{1}, +5)$            | $(\mathbf{3}, +1)$ | $(\mathbf{3}, +1)$ | $\sim \epsilon M$   |
| C     | $S'_4$ | $i\infty$           | $(\hat{\mathbf{1}}, +2) \oplus (\mathbf{1}, +3) \oplus (\hat{\mathbf{1}}, +4)$   | $(\mathbf{3}, +1)$ | $(\mathbf{3}, +1)$ | $\sim \epsilon^2 M$ |
| D     |        |                     | $(\hat{\mathbf{1}}', +2) \oplus (\mathbf{1}, +3) \oplus (\hat{\mathbf{1}}', +4)$ |                    |                    |                     |

Let's focus on this



Notation:  $\psi \sim (\rho, k_\psi)$

## An example: model C

Group  
 $\Gamma'_4 \simeq S'_4$

| $L$                                                                            | $E^c$              | $N^c$              | $\Delta M$          |
|--------------------------------------------------------------------------------|--------------------|--------------------|---------------------|
| $(\hat{\mathbf{1}}, +2) \oplus (\mathbf{1}, +3) \oplus (\hat{\mathbf{1}}, +4)$ | $(\mathbf{3}, +1)$ | $(\mathbf{3}, +1)$ | $\sim \epsilon^2 M$ |

$$\tau_{\text{sym}} \simeq i\infty$$

$$W = (Y_e)_{ij} L_i E_j^c H_d + (Y_D)_{i\kappa} L_i N_\kappa^c H_u + \frac{1}{2} (M_R)_{\kappa\rho} N_\kappa^c N_\rho^c$$



# An example: model C

Group  
 $\Gamma'_4 \simeq S'_4$

| $L$                                                                            | $E^c$              | $N^c$              | $\Delta M$          |
|--------------------------------------------------------------------------------|--------------------|--------------------|---------------------|
| $(\hat{\mathbf{1}}, +2) \oplus (\mathbf{1}, +3) \oplus (\hat{\mathbf{1}}, +4)$ | $(\mathbf{3}, +1)$ | $(\mathbf{3}, +1)$ | $\sim \epsilon^2 M$ |

$$\tau_{\text{sym}} \simeq i\infty$$

$$M_R = \Lambda \begin{pmatrix} \frac{2}{\sqrt{3}} (Y_2^{(2)})_1 & 0 & 0 \\ 0 & (Y_2^{(2)})_2 & -\frac{1}{\sqrt{3}} (Y_2^{(2)})_1 \\ 0 & -\frac{1}{\sqrt{3}} (Y_2^{(2)})_1 & (Y_2^{(2)})_2 \end{pmatrix}$$

$$Y_D = \begin{pmatrix} g_1 (Y_{\hat{\mathbf{3}}'}^{(3)})_1 & g_1 (Y_{\hat{\mathbf{3}}'}^{(3)})_3 & g_1 (Y_{\hat{\mathbf{3}}'}^{(3)})_2 \\ g_2 (Y_{\mathbf{3}}^{(4)})_1 & g_2 (Y_{\mathbf{3}}^{(4)})_3 & g_2 (Y_{\mathbf{3}}^{(4)})_2 \\ g_3 (Y_{\hat{\mathbf{3}}'}^{(5)})_1 & g_3 (Y_{\hat{\mathbf{3}}'}^{(5)})_3 & g_3 (Y_{\hat{\mathbf{3}}'}^{(5)})_2 \end{pmatrix}$$

$$\sim \begin{pmatrix} 1 & 0 & 0 \\ 0 & \epsilon^2 & 1 \\ 0 & 1 & \epsilon^2 \end{pmatrix}$$


$$\epsilon \sim q^{1/4} = e^{i\pi\tau/2}$$

$$Y_e = \begin{pmatrix} \alpha_1 (Y_{\hat{\mathbf{3}}'}^{(3)})_1 & \alpha_1 (Y_{\hat{\mathbf{3}}'}^{(3)})_3 & \alpha_1 (Y_{\hat{\mathbf{3}}'}^{(3)})_2 \\ \alpha_2 (Y_{\mathbf{3}}^{(4)})_1 & \alpha_2 (Y_{\mathbf{3}}^{(4)})_3 & \alpha_2 (Y_{\mathbf{3}}^{(4)})_2 \\ \alpha_3 (Y_{\hat{\mathbf{3}}'}^{(5)})_1 & \alpha_3 (Y_{\hat{\mathbf{3}}'}^{(5)})_3 & \alpha_3 (Y_{\hat{\mathbf{3}}'}^{(5)})_2 \end{pmatrix}$$

$$\sim \begin{pmatrix} \epsilon^3 & \epsilon^2 & 1 \\ \epsilon^2 & \epsilon & \epsilon^3 \\ \epsilon^3 & \epsilon^2 & 1 \end{pmatrix}$$

L-conserving for  $\epsilon \rightarrow 0$

## Pheno: model C (Inverted Ordering)





6 dimensionless  
parameters  
9 in total (includes  $\Lambda$  and  $\tau$ )



# Pheno: model C (Inverted Ordering)

$$\Theta_{\alpha j} \equiv v_u \hat{Y}_{\alpha j} / M_j.$$

$$\Theta_{\alpha}^2 \equiv \sum_{j=1}^3 |\Theta_{\alpha j}|^2 \quad \text{and} \quad \Theta^2 \equiv \sum_{\alpha=e, \mu, \tau} \Theta_{\alpha}^2.$$



$$M_{av} = \frac{\sum_i M_i}{3}$$

# Where to go from here?

Smallness of light neutrino masses

$$m_\nu \rightarrow 0 \text{ as } \epsilon \rightarrow 0$$

$$m_e/m_\mu \simeq \frac{1}{200}$$

$$m_\mu/m_\tau \simeq \frac{1}{17}$$

$$\epsilon \equiv |\tau - \tau_{sym}|$$

$$\Delta M \sim |\epsilon|^r \times M.$$

L-conserving for  $\epsilon \rightarrow 0$

CP conserving (in gCP mode)

$$\delta_{CP}, \alpha_{21}, \alpha_{31} \rightarrow 0 \text{ as } \epsilon \rightarrow 0$$



## Where to go from here?

$$\epsilon \equiv |\tau - \tau_{sym}|$$

Is spontaneous  
CP violation too  
much to ask?



$$\delta_{CP}, \alpha_{21}, \alpha_{31} \rightarrow 0 \text{ as } \epsilon \rightarrow 0$$



Thank you for the attention

Roma Tre  
Neutrino  Group

SCAN ME









# BACKUP SLIDES



# Fundamental domain of the benchmarks



# The numerics of the benchmarks

| Model (ordering)                      | A (NO)                          | B (NO)                          | C (IO)                          | D (NO)                          | D (IO)                       |
|---------------------------------------|---------------------------------|---------------------------------|---------------------------------|---------------------------------|------------------------------|
| $\text{Re } \tau$                     | $-0.472^{+0.017}_{-0.028}$      | $-0.475^{+0.036}_{-0.024}$      | all                             | all                             | all                          |
| $\text{Im } \tau$                     | $0.892^{+0.019}_{-0.042}$       | $0.912^{+0.015}_{-0.062}$       | $2.55^{+0.07}_{-0.04}$          | $2.31^{+0.25}_{-0.82}$          | $1.68^{+0.37}_{-0.24}$       |
| $\hat{\alpha}_2/\hat{\alpha}_1$       | $-0.286^{+0.674}_{-0.106}$      | $0.590^{+0.125}_{-0.083}$       | $-0.593^{+0.102}_{-0.078}$      | $0.31^{+1.06}_{-1.05}$          | $-0.058^{+0.348}_{-0.084}$   |
| $\hat{\alpha}_{3(,1)}/\hat{\alpha}_1$ | $-0.160^{+0.145}_{-0.481}$      | $-0.0502^{+0.0501}_{-0.1540}$   | $0.0339^{+0.0026}_{-0.0024}$    | $0.33^{+2.08}_{-0.12}$          | $-1.38^{+0.70}_{-0.85}$      |
| $\hat{\alpha}_{3,2}/\hat{\alpha}_1$   | $0.0202^{+0.0094}_{-0.0514}$    | $-0.0185^{+0.0037}_{-0.0062}$   | —                               | $0.20^{+4.25}_{-0.79}$          | $-0.71^{+2.30}_{-2.39}$      |
| $\hat{g}_2/\hat{g}_1$                 | $0.712^{+0.052}_{-0.076}$       | $0.219^{+0.011}_{-0.008}$       | $0.0799^{+0.0137}_{-0.0125}$    | $1.56^{+1.30}_{-1.19}$          | $2.55^{+1.29}_{-0.61}$       |
| $\hat{g}_{3(,1)}/\hat{g}_1$           | $0.491^{+0.196}_{-0.461}$       | $0.336^{+0.062}_{-0.161}$       | $13.2^{+1.1}_{-1.2}$            | $0.129^{+0.011}_{-0.073}$       | $0.0651^{+0.0172}_{-0.0163}$ |
| $\hat{g}_{3,2}/\hat{g}_1$             | $0.214^{+0.119}_{-0.551}$       | $0.289^{+0.018}_{-0.020}$       | —                               | $0.853^{+0.077}_{-0.734}$       | $0.798^{+0.224}_{-0.231}$    |
| $v_d \hat{\alpha}_1$ , GeV            | $0.594^{+0.070}_{-0.121}$       | $0.363^{+0.047}_{-0.055}$       | $0.176^{+0.005}_{-0.007}$       | $0.0689^{+0.0132}_{-0.0427}$    | $0.0384^{+0.0126}_{-0.0061}$ |
| $v_u^2 \hat{g}_1^2/\Lambda$ , eV      | $0.0328^{+0.0145}_{-0.0070}$    | $0.0568^{+0.0192}_{-0.0120}$    | $0.00032^{+0.00005}_{-0.00003}$ | $0.0184^{+0.0069}_{-0.0066}$    | $0.0120^{+0.0108}_{-0.0035}$ |
| $m_e/m_\mu$                           | $0.0048^{+0.0006}_{-0.0005}$    | $0.0047^{+0.0006}_{-0.0005}$    | $0.0048^{+0.0005}_{-0.0006}$    | $0.0047^{+0.0005}_{-0.0006}$    | $0.0048^{+0.0005}_{-0.0006}$ |
| $m_\mu/m_\tau$                        | $0.0560^{+0.0115}_{-0.0114}$    | $0.0571^{+0.0114}_{-0.0121}$    | $0.0577^{+0.0109}_{-0.0134}$    | $0.0550^{+0.0129}_{-0.0105}$    | $0.0572^{+0.0099}_{-0.0117}$ |
| $r$                                   | $0.0297^{+0.0022}_{-0.0021}$    | $0.0297^{+0.0023}_{-0.0021}$    | $0.0300^{+0.0023}_{-0.0021}$    | $0.0299^{+0.0021}_{-0.0023}$    | $0.0302^{+0.0020}_{-0.0023}$ |
| $\sin^2 \theta_{12}$                  | $0.307^{+0.034}_{-0.030}$       | $0.308^{+0.032}_{-0.030}$       | $0.312^{+0.019}_{-0.018}$       | $0.310^{+0.030}_{-0.034}$       | $0.308^{+0.032}_{-0.034}$    |
| $\sin^2 \theta_{13}$                  | $0.0220^{+0.0017}_{-0.0014}$    | $0.0221^{+0.0014}_{-0.0014}$    | $0.0222^{+0.0016}_{-0.0014}$    | $0.0222^{+0.0016}_{-0.0016}$    | $0.0224^{+0.0013}_{-0.0015}$ |
| $\sin^2 \theta_{23}$                  | $0.506^{+0.067}_{-0.065}$       | $0.507^{+0.065}_{-0.062}$       | $0.519^{+0.049}_{-0.058}$       | $0.512^{+0.070}_{-0.065}$       | $0.507^{+0.070}_{-0.030}$    |
| $m_1$ , eV                            | $< 10^{-4}$                     | $< 10^{-4}$                     | $0.0491^{+0.0002}_{-0.0002}$    | $0.00267^{+0.00048}_{-0.00132}$ | $0.0620^{+0.0025}_{-0.0032}$ |
| $m_2$ , eV                            | $0.00864^{+0.00028}_{-0.00029}$ | $0.00865^{+0.00029}_{-0.00028}$ | $0.0499^{+0.0001}_{-0.0002}$    | $0.00907^{+0.00028}_{-0.00044}$ | $0.0626^{+0.0024}_{-0.0032}$ |
| $m_3$ , eV                            | $0.0501^{+0.0001}_{-0.0002}$    | $0.0501^{+0.0002}_{-0.0002}$    | $< 10^{-4}$                     | $0.0502^{+0.0002}_{-0.0002}$    | $0.0379^{+0.0038}_{-0.0054}$ |
| $\Sigma_i m_i$ , eV                   | $0.0588^{+0.0001}_{-0.0001}$    | $0.0588^{+0.0001}_{-0.0001}$    | $0.0990^{+0.0003}_{-0.0004}$    | $0.0619^{+0.0006}_{-0.0017}$    | $0.163^{+0.009}_{-0.012}$    |
| $m_{\beta\beta}$ , meV                | $1.49^{+0.31}_{-0.28}$          | $1.49^{+0.27}_{-0.28}$          | $17.8^{+1.7}_{-1.8}$            | $2.06^{+0.98}_{-0.22}$          | $60.0^{+2.4}_{-3.1}$         |
| $\delta$                              | $\simeq 0, \pi$                 | $\simeq 0$                      | $\simeq 0$                      | $\simeq \pi$                    | $\simeq 0, \pi$              |
| $\alpha_{21(23)}$                     | $\simeq \pi$                    | $\simeq \pi$                    | $\simeq 0$                      | $\simeq \pi$                    | $\simeq 0$                   |
| $\alpha_{31}$                         | —                               | —                               | —                               | $\simeq \pi$                    | $\simeq \pi$                 |
| $ \epsilon(\tau) $                    | $0.0218^{+0.0047}_{-0.0046}$    | $0.0292^{+0.0076}_{-0.0067}$    | $0.0182^{+0.0013}_{-0.0018}$    | $0.0267^{+0.0700}_{-0.0087}$    | $0.0713^{+0.0320}_{-0.0312}$ |
| $(m_e/m_\tau)/ \epsilon ^{2,3}$       | $0.563^{+0.167}_{-0.111}$       | $0.317^{+0.123}_{-0.088}$       | $45.8^{+3.1}_{-3.1}$            | $14.0^{+26.0}_{-13.7}$          | $0.76^{+2.85}_{-0.50}$       |
| $(m_\mu/m_\tau)/ \epsilon $           | $2.57^{+0.05}_{-0.08}$          | $1.96^{+0.20}_{-0.24}$          | $3.16^{+0.39}_{-0.52}$          | $2.06^{+0.96}_{-1.53}$          | $0.801^{+0.517}_{-0.200}$    |
| $\min N\sigma$                        | 0.412                           | 0.411                           | 0.548                           | 0.488                           | 0.362                        |
| max par. ratio                        | 49.6                            | 54.1                            | 165                             | 12.2                            | 39.1                         |



# Potential shortcomings

## Bottom-up problems

Kähler  
potential

Normalisation of the  
modular forms


Modulus  
stabilization



SUSY-dependent theory





Three “copies”, different masses



non-abelian discrete symmetries

$$S_3 \quad A_4 \quad S_4 \quad A_5 \dots\dots$$

$$\mathcal{W}_{Yukawa} \supset \frac{\alpha}{\Lambda} E^c (L\phi_i)_1 H_d$$



Isidor I. Rabi

$$U_{\text{TBM}} = \begin{pmatrix} \sqrt{\frac{2}{3}} & \frac{1}{\sqrt{3}} & 0 \\ -\frac{1}{\sqrt{6}} & \frac{1}{\sqrt{3}} & -\frac{1}{\sqrt{2}} \\ -\frac{1}{\sqrt{6}} & \frac{1}{\sqrt{3}} & \frac{1}{\sqrt{2}} \end{pmatrix}$$

Tri-bimaximal  
compatible with data  
until 2012

$$\theta_{13} = 0 \quad \theta_{12} \simeq 35^\circ \quad \theta_{23} = 45^\circ$$

Not zero

F. P. An *et al.*, “Observation of electron-antineutrino disappearance at daya bay,” *Phys. Rev. Lett.* **108** (Apr, 2012)



## Shortcomings of the traditional approach

EFT with scalar “flavons”  $\phi_i$

✗  $U_{\text{PMNS}} = U_{\text{TBM}}^0 + \dots \text{corrections}$

$\downarrow$   
 $\theta_{13} = 0$   
 Automatic

$\downarrow$   
 $\theta_{13} \approx 8.5^\circ ?$

✗

$\delta\theta_{23}, \delta\theta_{12} \approx 8.5^\circ$

✗  $V(\phi_i) \rightarrow \text{Mess!}$

# The Modular symmetry approach

## Modular-invariant SUSY action

$$\mathcal{S} = \int d^4x \int d^2\theta d^2\bar{\theta} K(\Phi, \bar{\Phi}) + \left[ \int d^4x \int d^2\theta \mathcal{W}(\Phi) + \text{h.c.} \right]$$

Kähler potential

Superpotential

$$\sigma \equiv \Lambda_\tau \tau$$

- ▶ Gives the kinetic terms after the modulus acquires a VEV
- ▶ A minimalistic form is chosen

- ▶ Holomorphic function of superfields
- ▶ Encodes the Higgs Yukawa interactions

The superfields transform as:

$$\begin{cases} \tau \rightarrow \gamma(\tau) = \frac{a\tau + b}{c\tau + d} \\ \varphi^{(I)} \rightarrow (c\tau + d)^{-k_I} \rho^{(I)}(\gamma) \varphi^{(I)} \end{cases}, \quad \text{with } \gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma_N$$

Action is invariant if,  
under  $\Gamma_N$ :

$$\begin{cases} \mathcal{W}(\Phi) \rightarrow \mathcal{W}(\Phi) \\ K(\Phi, \bar{\Phi}) \rightarrow \underbrace{K(\Phi, \bar{\Phi}) + f(\Phi) + f(\bar{\Phi})}_{\text{Kähler transformation}} \end{cases}$$

$\theta, \bar{\theta}$  Grassmann spinor coordinates

$\Phi = (\tau, \varphi)$  Chiral superfields

$\varphi$  Usual matter supermultiplets

$\rho(\gamma)$  Unitary representation of  $\Gamma_N$



# The Kähler potential...

- ▶ Minimalist choice for the Kähler

$$K(\Phi, \bar{\Phi}) = -h\Lambda_\tau^2 \log(-i\tau + i\bar{\tau}) + \sum_I (-i\tau + i\bar{\tau})^{-k_I} |\varphi^{(I)}|^2$$

$\Lambda_\tau \equiv$  dimensions of mass

Satisfies  $K(\Phi, \bar{\Phi}) \rightarrow K(\Phi, \bar{\Phi}) + f(\Phi) + f(\bar{\Phi})$  under  $\Gamma_N$   $h \equiv$  positive constant

- ▶ In a bottom-up approach, this is just a choice

- ▶ Corrections of the Kähler potential can spoil the predictivity of the model

M.-C. Chen, S. Ramos-Sánchez, and M. Ratz, “A note on the predictions of models with modular flavor symmetries,” *Physics Letters B* **801** (Feb, 2020) 135153.

- ▶ This question is an open one

# The Modular symmetry approach

## The group generators

► Finite modular group can be defined:  $\Gamma_N \equiv \bar{\Gamma}/\bar{\Gamma}(N)$

$$\Gamma(N) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \text{SL}(2, \mathbb{Z}) \mid \begin{pmatrix} a & b \\ c & d \end{pmatrix} \equiv \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \pmod{N} \right\}$$

subgroups of  $\Gamma$   $N=1,2,3\dots$  called “level”

$$\begin{aligned} \bar{\Gamma} &\equiv \Gamma/\{\pm \mathbb{I}\} \\ \bar{\Gamma}(N) &\equiv \Gamma(N)/\{\pm \mathbb{I}\} \end{aligned}$$

► Generators S and T of the modular group  $\Gamma_N$

$$\begin{array}{cc} \text{S} & \text{T} \\ \tau \rightarrow -\frac{1}{\tau} & \tau \rightarrow \tau + 1 \end{array} \quad S^2 = T^N = (ST)^3 = \mathbb{I}$$

$$S = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \quad T = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$$

►  $S_3$  Generators S and T satisfy:

$$S^2 = T^2 = (ST)^3 = \mathbb{I}$$

$$\rho(S) = \frac{1}{2} \begin{pmatrix} -1 & -\sqrt{3} \\ \sqrt{3} & 1 \end{pmatrix}, \quad \rho(T) = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.$$

$$(\rho(S))^2 = \mathbb{I}, \quad (\rho(S)\rho(T))^3 = \mathbb{I}, \quad (\rho(T))^2 = \mathbb{I},$$



# Backup slides

## Numerical procedure

- ▶ Define a “figure of merit”, i.e. chi-square for every set of parameters  $l(p_i) \equiv \sqrt{\chi^2(p_i)}$

$$\chi^2(p_i) = \sum_{j=1}^6 \left( \frac{q_j(p_i) - q_j^{\text{b-f}}}{\sigma_j} \right)^2$$

$$p_i = \{\tau, \beta/\alpha, \gamma/\alpha, \dots, g'/g, g_p/g, \dots\}$$


- ▶ Define a “potential” with a given temperature  $T$  and a threshold

$$V(p_i) = \begin{cases} l(p_i) & , \quad l(p_i) \leq l_{\text{max}} \\ +\infty & , \quad \text{otherwise} \end{cases}$$

$$q_j = \{\sin^2 \theta_{12}, \sin^2 \theta_{13}, \sin^2 \theta_{23}, m_e/m_\mu, m_\mu/m_\tau, r\}$$

P. P. Novichkov, J. T. Penedo, S. T. Petcov, and A. V. Titov,  
“Modular  $S_4$  models of lepton masses and mixing,” (2019)

- ▶ At iteration “t”, generate a new point from a Gaussian centred on the previous one



- ▶ Accept the new point with a probability given by:

$$P_\alpha = \min[1, \exp(V(p_i^{(t)}) - V(p'_i))/T]$$

# Non-standard interactions

## tests of modulus couplings

G-J. Ding, FF,  
2003.13448

non standard neutrino  
interactions

$$\mathcal{L} = i \sum_{f=e,e^c,\nu} \bar{f} \not{\partial} f + \frac{1}{2} \partial_\mu \varphi_\alpha \partial^\mu \varphi_\alpha - \frac{1}{2} M_\alpha^2 \varphi_\alpha^2$$

$$- (m_e + Z_\alpha^e \varphi_\alpha) e^c e - \frac{1}{2} \nu (m_\nu + Z_\alpha^\nu \varphi_\alpha) \nu + h.c. + \dots$$


$$\tau = \langle \tau \rangle + \frac{\varphi_u + i \varphi_v}{\sqrt{2}}$$

in medium with non-zero  
electron number density

small, unless the modulus is very light

$$\delta m_\nu(0) = -n_e^0 \frac{\text{Re}(Z^e) Z^\nu}{M^2(R)},$$

in the sun:



from Feruglio's slides at Mod. Symmetry Bethe Workshop