



Contribution ID: 171

Type: **not specified**

Lack of propagating degrees of freedom in degenerate $f(R)$ models

Tuesday 27 May 2025 18:40 (20 minutes)

In the context of $f(R)$ gravity, as well as other extended theories of gravity, the correct counting of globally well-defined linear dynamical modes (i.e. gravitational-wave polarisations) has recently drawn a vivid interest. In this talk, we present a consistent approach shedding light on such issues for both so-called degenerate and non-degenerate $f(R)$ models embedded in Minkowski and de Sitter backgrounds. We find that the linearised spectrum of degenerate models on these backgrounds is empty, lacking both the graviton and scalaron modes which appear in generic non-degenerate models. Our work generalises previous results in the literature applicable only to the specific (degenerate) model $f(R) = \alpha R^2$; in fact, we find that the same pathologies discovered therein emerge for all choices of $f(R)$ belonging to the wide class of degenerate models.

Primary author: CASADO-TURRIÓN, Adrián (Center for Physical Sciences and Technology (FTMC, Lithuania))

Co-authors: DOBADO, Antonio (Complutense University of Madrid); DE LA CRUZ-DOMBRIZ, Álvaro (University of Salamanca and University of Cape Town)

Presenter: CASADO-TURRIÓN, Adrián (Center for Physical Sciences and Technology (FTMC, Lithuania))

Session Classification: Formal