



Istituto Nazionale di Fisica Nucleare  
Laboratori Nazionali di Frascati

# Baryogenesis via bubble collisions

**Kristjan Mürsepp**, *in collaboration with*

**Martina Cataldi** (UHH, DESY) and **Miguel Vanvlasselaer** (Brussels U, IIHE)

Laboratori Nazionali di Frascati (INFN - LNF), Frascati, Italy

Keemilise-Bioloogilise Füüsika Instituut (KBFI), Tallinn, Estonia

PLANCK 2025,

*Centro Culturale Altinate San Gaetano*

**29.05.2025**



Keemilise ja  
Bioloogilise Füüsika Instituut

National Institute of Chemical Physics and Biophysics




# The asymmetry of the Universe

The Universe contains more matter than antimatter as evidenced by the structure around us.

This can be observed in two equivalent ways:


These values are inferred independently from

1. Enhancement of the odd peaks in CMB
2. Abundances of D,  $^3\text{He}$  during BBN



Credit: ESA

$$\eta \equiv \left. \frac{n_B - n_{\bar{B}}}{n_\gamma} \right|_0 = (6.21 \pm 0.16) \times 10^{-10},$$
$$Y_{\Delta B} \equiv \left. \frac{n_B - n_{\bar{B}}}{s} \right|_0 = (8.75 \pm 0.23) \times 10^{-11}$$



Credit: Wikipedia



# Sakharov conditions

The observed baryon asymmetry should be created dynamically after inflation:

This can be achieved by the 3 Sakharov conditions:

1. Baryon number violation
2. C- and CP- violation
3. Departure from thermal equilibrium



Credit: Wikipedia



## Asymmetrizing within the SM

Can we satisfy Sakharov conditions in the SM...? **Not really!**

$$\Delta B = \Delta L = \pm 3.$$

1. **Sphalerons** conserve B-L but violate B+L  
with *unsuppressed* decay rate at **high temperatures**. **OK!**
- 2.1 Weak interactions **violate C maximally**. **OK!**
- 2.2 The CP-violation from the CKM matrix however is  $\sim 10^{-20}$ . **NOT ENOUGH!**
3. EWSB does not give a first order phase transition – Higgs is too heavy! **NOT OK!**

**Look for explanation in connection with other unsolved problems of the SM.**



# Neutrinos to the rescue

Baryogenesis can be nicely connected to an explanation of neutrino masses.

1. Light neutrino masses from a heavy right-handed neutrino  $\rightarrow$  *seesaw mechanism*
2. *CP violating* decays of RHN are *out of equilibrium* and produce  $\Delta L$
3.  $\Delta L$  is converted into  $\Delta B$  by *sphalerons*.

Features of the THERMAL LEPTOGENESIS scenario:

1. RHN has to be very heavy  $m_N > 10^9 \text{ GeV} \Rightarrow$  High reheating temperatures needed
2. Not many clear experimental signatures.
3. CP-asymmetry in the DECAY of *on-shell* RHN.

LINK WITH PHASE TRANSITIONS.

We use the ideas first outlined in

M.Cataldi and B.Shakya, *JCAP* **11** (2024) 047

Artwork by Sandbox Studio, Chicago with Ana Kova





# The boiling Universe

The key observables that we are interested in:

1. **Energy released by the PT**

$$\Delta V \equiv V_{\langle\phi\rangle=0} - V_{\langle\phi\rangle=v} = c_V v^4$$

2. **Inverse duration of the PT**

$$\beta(T) = T \frac{d}{dT} \left( \frac{S_3}{T} \right)$$

3. **Lorentz boost factor of the bubble**

$$\gamma_w = 1/\sqrt{1 - v_w^2}.$$


4. **Bubble wall thickness**

$$l_w = l_{w0}/\gamma_w \text{ with } l_{w0} \sim \mathcal{O}(v_\phi^{-1})$$


5. **Bubble radius at collision**

$$R_{\text{coll}} \approx \frac{(8\pi)^{1/3} v_w}{H(T_{\text{reh}}) \beta(T_{\text{reh}})}.$$

**Strong phase transitions often feature gravitational wave signals!**



# Particles popping from bubbles



For that *runaway bubbles* are needed.

We need to limit the *friction* on the bubble walls

$$\mathcal{P}_{\text{LO}} \approx \frac{1}{24} m^2 T^2$$

$$\mathcal{P}_{\text{NLO}} \sim g^2 \gamma_w m_V T^3$$

Dorsch, Huber, Konstandin,

*JCAP* **12** (2018) 034

Bödeker, Moore

*JCAP* **05**, *JCAP* **05** (2009) 009

Bödeker, Moore *JCAP* **05** (2017) 025

Gouttenoire, Jinno, Sala *JHEP* **05** (2022) 004

Ai, *JCAP* **10** (2023) 052,

Long, Turner, *JCAP* **11** (2024) 024

Maximum energy for particle production  $E \sim \gamma v$

$$\gamma_w^{\text{coll}} \sim \frac{2\sqrt{10} M_{\text{pl}} T_{\text{nuc}} (8\pi)^{1/3} v_w}{\pi \sqrt{g_*} \beta T_{\text{reh}}^2} \approx 5.9 \frac{M_{\text{pl}} T_{\text{nuc}} v_w}{\sqrt{g_*} \beta T_{\text{reh}}^2}.$$

Compared to bubble – plasma collision :  $v_w$  does not have to be related to the heavy particle mass!



# Zooming in on the bubble collisions

The number of particles produced per unit area:

$$\frac{N}{A} = \frac{1}{2\pi^2} \int_{p_{\min}^2}^{p_{\max}^2} dp^2 f(p^2) \text{Im}[\tilde{\Gamma}^{(2)}(p^2)] .$$

A. Falkowski, J.M.No *JHEP* **02** (2013) 034

The imaginary part of the 1PI 2-point Green function

$$\text{Im}[\tilde{\Gamma}^{(2)}(p^2)] = \frac{1}{2} \sum_k \int d\Pi_k |\bar{\mathcal{M}}(\phi_p^* \rightarrow k)|^2$$

R. Watkins and L. M. Widrow, *Nuc. Phys B* **02** (1992) 347

The Fourier transform has been computed numerically:

$$f_{\text{elastic}}(p^2) = f_{\text{PE}}(p^2) + \frac{v_\phi^2 L_p^2}{15m_t^2} \exp\left(\frac{-(p^2 - m_t^2 + 12m_t/L_p)^2}{440 m_t^2/L_p^2}\right) \quad (\text{elastic collisions})$$

$$f_{\text{inelastic}}(p^2) = f_{\text{PE}}(p^2) + \frac{v_\phi^2 L_p^2}{4m_f^2} \exp\left(\frac{-(p^2 - m_f^2 + 31m_f/L_p)^2}{650 m_f^2/L_p^2}\right) \quad (\text{inelastic collisions})$$

Number of particles produced

H. Mansour, B.Shakya. *Phys.Rev.D* **111** (2025) 2

For very heavy particles only the first part is important

$$f_{\text{PE}}(p^2) = \frac{16v_\phi^2}{p^4} \text{Log} \left[ \frac{2(1/l_w)^2 - p^2 + 2(1/l_w) \sqrt{(1/l_w)^2 - p^2}}{p^2} \right]$$

$$\frac{n}{s} = \frac{1}{s(T_{\text{reh}})} \overbrace{\left. \frac{N}{A} \right|_N}^{\text{production per surface}} \times \overbrace{\frac{3}{2R_{\text{coll}}}}^{\text{diffusion}}$$



# Introducing the heavyweights

We consider the simplified Lagrangian

with the hierarchy

$$\mathcal{L} = Y\phi P_R \bar{N} \chi + \frac{1}{2} M_N N \bar{N} + \frac{1}{2} m_\chi \chi \bar{\chi} + \sum_\alpha y_\alpha P_R N (\tilde{H} \bar{L}_\alpha) - V(\phi, T)$$

$$m_N \gg m_\chi \gg T_{\text{reh}} \sim v \gg v_{\text{EW}}.$$

The yield of the heavy particles produced is given by

$$Y_N^{\text{BC}} \simeq 0.012 N |Y|^2 \frac{\beta}{v_w} \left( \frac{\pi^2 \alpha}{30(1+\alpha) g_* c_V} \right)^{1/4} \frac{v}{M_{\text{pl}}} \log \left( \frac{2\gamma_w v}{m_\chi + m_N} \right)$$

To avoid **backreaction** one needs

which gives a rough constraint

$$\rho_N^{\text{BC}} \approx 0.03 N |Y|^2 v^5 \frac{T_{\text{nuc}}}{T_{\text{reh}}^2} \sim 0.03 N |Y|^2 v^3 T_{\text{nuc}} \ll \Delta V = c_V v^4$$



$$0.03 N \times |Y|^2 \frac{T_{\text{nuc}}}{v} \ll c_V$$



# Boosted light(ning) fast particles

Boosted SM particles can also be produced directly via *off-shell* N

$$\Gamma_{\phi^* \rightarrow HL\chi}(p^2) = \frac{2}{(2\pi)^3} \frac{|y|^2 |Y|^2}{32\sqrt{p^6}} \int_{s_{12}^{\min}}^{s_{12}^{\max}} \int_{s_{23}^{\min}}^{s_{23}^{\max}} \frac{m_N^2 (s_{23} - m_L^2 - m_\chi^2)}{(s_{12} - m_N^2)^2 + m_N^2 \Gamma_N^2} ds_{12} ds_{23}.$$




The production is always dominated by the mN resonance!

Low energies	$p \ll m_N :$	$\Gamma_{\phi \rightarrow HL\chi} \simeq 2 \frac{ y ^2  Y ^2}{1536\pi^3} \frac{p^3}{m_N^2}$
High energies	$p \gg m_N :$	$\Gamma_{\phi \rightarrow HL\chi} \simeq 2 \frac{ y ^2  Y ^2 p}{512\pi^2} \frac{m_N}{\Gamma_N}$



# CP-violation in the *production*

We generalize the Lagrangian to include flavours



$$\mathcal{L} = \sum_{iI} Y_{iI} \phi \bar{N}_i P_R \chi_I + \sum_{i\alpha} y_{i\alpha} P_R N_i (\tilde{H} \bar{L}_\alpha) + \sum_i M_i^N \bar{N}_i N_i + \sum_I M_I^\chi \bar{\chi}_I \chi_I - V(\phi, T) + h.c.$$

**Equal and opposite** asymmetries produced, but **SEPARATED** into two sectors

$$\begin{aligned} \epsilon_{iI} &\equiv \frac{|\mathcal{M}_{\phi \rightarrow N_i \chi_I^c}|^2 - |\mathcal{M}_{\phi \rightarrow N_i^c \chi_I}|^2}{\sum_{iI} |\mathcal{M}_{iI}|^2 + |\mathcal{M}_{\bar{i}\bar{I}}|^2} \\ &= \frac{2 \sum_{j,J} \text{Im}(Y_{iI} Y_{iJ}^* Y_{jJ} Y_{jI}^*) \text{Im} f_{ij}^{(\chi\phi)}}{\sum_{i,I} |Y_{iI}|^2} + \frac{2 \sum_{\alpha,j} \text{Im}(Y_{iI} y_{i\alpha}^* y_{j\alpha} Y_{jI}^*) \text{Im} f_{ij}^{(HL)}}{\sum_{i,I} |Y_{iI}|^2} \end{aligned}$$

$$n_{N_i} - n_{\bar{N}_i} = n_{\Delta N_i}^i \approx \sum_I \epsilon^{iI} n_{N_i} \quad n_{\chi_I} - n_{\bar{\chi}_I} = n_{\Delta \chi_I} \approx - \sum_i \epsilon^{iI} n_{\chi_I}$$

$$n_{\Delta N} + n_{\Delta \chi} = 0.$$

**N** talks to the SM => **transmits its asymmetry to the SM**

**chi** talks to a light dark sector => its asymmetry is **secluded** from the SM.



# Combining CP violation from *production* and *decay*

$$\begin{aligned} \frac{n_{L_\alpha} - n_{\bar{L}_\alpha}}{s(T_{\text{nuc}})} &\approx \frac{1}{s(T_{\text{nuc}})} \left( \sum_i \epsilon^{i\alpha} n_{N_i} + \sum_{Ii} \epsilon^{iI} n_{N_i} \right) \text{Br}[N_i \rightarrow HL_\alpha] \\ &\approx \frac{1}{32\pi} \sum_i \frac{n_{N_i}}{s(T_{\text{nuc}})} \left( \frac{\sum_{I,j \neq i} \text{Im}[y_{j\alpha}^* y_{\alpha i} Y_{iI} Y_{jI}^*] \frac{m_j m_i}{m_j^2 - m_i^2}}{\sum_{i,\beta} |y_{i\beta}|^2} + \right. \\ &\quad \left. 2 \times \frac{\sum_{j \neq i,\beta} \text{Im}[y_{i\beta} y_{\beta j}^* Y_{jI}^* Y_{iI}] \frac{m_i m_j}{m_j^2 - m_i^2}}{\sum_{i,I} |Y_{iI}|^2} \right) \text{Br}[N_i \rightarrow HL_\alpha], \end{aligned}$$

← CP-violation from the *production* of N

← Fraction of asymmetry transferred to SM




CP violation from the *decay* of N

Bubble wall plasma contribution subdominant due to

$$m_N \gg m_\chi \gg T_{\text{reh}} \sim v \gg v_{\text{EW}}.$$



# Production via *off-shell* $N$



$$\epsilon \equiv \frac{|\mathcal{M}|^2_{\phi \rightarrow \bar{\chi} \tilde{H} L} - |\mathcal{M}|^2_{\phi \rightarrow \chi H \bar{L}}}{|\mathcal{M}|^2_{\phi \rightarrow \bar{\chi} \tilde{H} L} + |\mathcal{M}|^2_{\phi \rightarrow \chi H \bar{L}}}.$$

$$(\Gamma^{\epsilon}_{\phi^* \rightarrow HL\chi})_{\alpha I} = \frac{1}{(2\pi)^3} \frac{1}{32p^3} \int_{s_{12}^{\min}}^{s_{12}^{\max}} \int_{s_{23}^{\min}}^{s_{23}^{\max}} |\mathcal{M}_0^{\alpha I}(s_{12}, s_{23})|^2 \epsilon_{\alpha I}(s_{12}) ds_{12} ds_{23}$$

The rate of production of the asymmetry

The asymmetry produced as a result:

$$\frac{N_{\Delta L}}{A} \Big|_{\phi^* \rightarrow \chi HL} \approx \int \frac{dp_z d\omega}{(2\pi)^2} \Gamma_{\phi^* \rightarrow \chi HL}(p) |\phi(p^2)|^2 \Rightarrow n_{\Delta L} \approx \frac{3\beta H}{2} \times \frac{N_{\Delta L}}{A} \Big|_{\phi^* \rightarrow \chi HL}$$



# Two special cases of the *off-shell* production

We focus on 2 generations of N, and assume  $m_1 \ll m_2$  and  $m_2 \gg \gamma v$



a. When  $\gamma v \gg m_1$ :

1. N1 resonance dominates the 1- $\rightarrow$ 3 production
2. We recover the CP-violation from the production and decay of an *on-shell* N1.

b. When  $\gamma v < m_1$ :

1. CP-violation can still occur via *off-shell* production

$$\left. \frac{N_{\Delta L}}{A} \right|_{\phi \rightarrow \chi HL} \approx 10^{-3} \frac{N \times C}{16\pi^6} (|y|^4 |Y|^2 \frac{m_1}{m_2} + \frac{1}{2} |y|^2 |Y|^4) \frac{p_{\max}^4 v^2}{m_1^3 m_2}$$





# WASH-OUTS



credit: Walmart

Thermalization much faster than the inverse decay

$$\tau_{HL \rightarrow N} \sim \frac{1}{\frac{|y|^2}{8\pi} \frac{T m_N^2}{E_{L,\text{initial}}^2}} \text{Exp} \left[ \frac{m_N^2}{E_{L,\text{initial}} T} \right] > \tau_{\text{therm}} \sim \frac{64\pi^3}{g^4} \frac{E_{L,\text{initial}}}{T^2}$$

Lepton asymmetry produced is suppressed by

$$Y_{\Delta L}^{\text{fin}} \approx \left( \Pi_i W_i \right) Y_{\Delta L}^{\text{init}},$$

$$W_{HL \rightarrow \phi \chi} \approx \text{Exp} \left[ - \frac{|y|^2 |Y|^2 M_{\text{pl}} T_{\text{reh}}}{0.3 \times 64\pi^5 m_N^2} \right].$$

$$W_{HL \rightarrow N} \approx \text{Exp} \left[ - \frac{2}{g_*^{1/2}} \left( \frac{M_{\text{pl}}}{m_N} \right) \left( \frac{|\sum_{\alpha} y_{\alpha 1}|^2}{8\pi} \right) \times \Gamma \left[ \frac{7}{2}, \frac{m_N}{T_{\text{reh}}} \right] \right].$$

Washout can be easily avoided for  $m_N/T \gg 10$



# Two birds with one stone : a tale of *cogenesis*

To explain the *coincidence problem* we slightly extend the Lagrangian


$$\mathcal{L} \supset y_1 \bar{\chi}(\tilde{\chi}\tilde{\phi}) + y_2 \phi \bar{\chi}\chi + \lambda_{\tilde{\phi}H}|H|^2\tilde{\phi}^2 - \frac{1}{2}m_{\tilde{\chi}}\tilde{\chi}\tilde{\bar{\chi}} + h.c.,$$

Only connection to the SM



Sizeable Higgs portal, (kinetic mixing)

Avoids washout from DS decays to SM



# Light from heavy: inverse seesaw mechanism

Introduce **lepton number violating** terms

$$\mathcal{L} \supset \sum_I \lambda_{N,R} \phi N_{R,I} \bar{N}_{R,I}^c + \lambda_{N,L} \phi N_{L,I} \bar{N}_{L,I}^c + \lambda_{\chi,R} \phi \chi_{R,I} \bar{\chi}_{R,I}^c + \lambda_{\chi,L} \phi \chi_{L,I} \bar{\chi}_{L,I}^c.$$

This results in the effective Weinberg operator

$$\mathcal{O}_{\text{Weinberg}} = \sum_{I,\alpha,\beta} \frac{y_{\alpha I} y_{\beta I}^* (\bar{L}_{\alpha}^c H)(L_{\beta} H) \lambda_{N,R} v}{m_N^2},$$

Giving a mass to the heaviest light neutrino



$$\text{Max}[m_{\nu}] \sim \text{Max} \left[ \sum_I |y_{\alpha I}|^2 \right] \frac{v_{EW}^2 \lambda_{N,R} v}{m_N^2}.$$



and the neutrino mass matrix

$$M = \begin{pmatrix} 0 & 0 & y v_H & 0 & 0 \\ 0 & \lambda_{L,R} v & m_N & 0 & Y v \\ y^T v_H & m_N^T & \lambda_{N,R} v & 0 & 0 \\ 0 & 0 & 0 & \lambda_{\chi,L} v & m_{\chi} \\ 0 & Y^T v & 0 & m_{\chi}^T & \lambda_{\chi R} v \end{pmatrix}$$

$$W_{H^c L \rightarrow H L^c} \approx \text{Exp} \left[ - \frac{6}{2\pi^3 g_{\star}^{1/2}} \frac{M_{\text{pl}} T_{\text{reh}} \sum m_{\nu_i}^2}{v_{EW}^4} \right].$$



# Parameter space for *on-shell* and *off-shell* production




# Gravitational wave signal

Since we have  $\gamma \gg 1$ , we will use the *bulk flow* model, T.Konstandin, *JCAP* **03** (2018) 047, R.Jinno, M. Takimoto, *JCAP* **01** (2019) 060

$$h^2 \Omega_{\text{GW}}^{\text{today}} = h^2 \Omega_{\text{peak}} S(f, f_{\text{peak}}) \quad S(f, f_{\text{peak}}) = \frac{(a+b) f_{\text{peak}}^b f^a}{b f_{\text{peak}}^{(a+b)} + a f^{(a+b)}}, \quad (a, b) \approx (0.9, 2.1),$$

energy mostly in the **sound waves**





Fluid sources GW long after collision

The energy density and the peak frequency are given by

$$h^2 \Omega_{\text{peak}} \approx 1.06 \times 10^{-6} \left( \frac{H_{\text{reh}}}{\beta} \right)^2 \left( \frac{\alpha \kappa}{1 + \alpha} \right)^2 \left( \frac{100}{g_\star} \right)^{1/3} \quad \text{and ,}$$
$$f_{\text{peak}} \approx 2.12 \times 10^{-3} \left( \frac{\beta}{H_{\text{reh}}} \right) \left( \frac{T_{\text{reh}}}{100 \text{ GeV}} \right) \left( \frac{100}{g_\star} \right)^{-1/6} \quad \text{mHz .}$$



# Gravitational wave signals for the Einstein Telescope (ET)



# Conclusions

- I. Bubble collisions from phase transitions allow to create very heavy particles
- II. The same mechanism can also entail CP violation in the production
- III. We have applied this observation to a concrete realization of baryogenesis
- IV. We computed CP violation both from the on-shell and off-shell production
- V. We start with zero asymmetry  $\Rightarrow$  the asymmetry is separated into two different sectors.
- VI. Consequently, we also attempt to explain the DM abundance via *cogenesis*.
- VII. Adding lepton violating terms can be used to also explain the neutrino masses

**Thank you for attention!**