


Padova,
14th September 2022

Supernovae as axion factories: the latest developments

Pierluca Carenza
OKC, Stockholm University

Core-Collapse Supernovae


For massive stars ($M > 8M_{\odot}$) the nuclear fusion produces heavy elements in an onion structure and a degenerate iron core



Iron in the core cannot be burnt and the star starts to collapse

Orders of magnitude for SNe

The SN core is an extreme environment



1000x



density

$$10^{14} \text{ g cm}^{-3}$$

temperature


$$30 \text{ MeV}$$

magnetic field

$$10^{15} \text{ G}$$

SN1987A: neutrino signal

From the few $\bar{\nu}_e p \rightarrow n e^+$ events of SN 1987A we know that...




$\sim 10^{53}$ erg emitted as neutrinos with energy $\sim O(15 \text{ MeV})$ in ~ 10 s

The energy-loss argument

G. Raffelt, Lect. Notes Phys. **741** (2008)

Stars produce axions which escape, draining energy from the core



Axions affect strongly the SN neutrino burst if


$$L_a > L_\nu = 2 \times 10^{52} \text{ erg s}^{-1}$$

Axion production in nuclear processes

Axion-nucleon bremsstrahlung in SNe

M. S. Turner, Phys. Rev. Lett. **60** (1988)

SN axions are produced by nucleon-axion bremsstrahlung



where we have to include detailed nuclear physics and many body effects

Pion-axion conversion in SNe

PC, B. Fore *et al.*, Phys. Rev. Lett. **126** (2021) no.7, 071102

SN axions are produced by pion-axion conversion




This is the leading axion production process in a SN despite the small density of pions ($\mathcal{O}(1\%)$)!!

Flux from pion-axion conversion

T. Fischer, PC *et al.* [[arXiv:2108.13726 \[hep-ph\]](#)].

The harder spectrum is due to the pion rest mass



Comparison of axion fluxes at $t_{\text{pb}} = 1 \text{ s}$

Consequences on the SN cooling

The SN cooling is accelerated by this new process, then the SN1987A bound is a factor 2 stronger

ρ		$\bar{g}_{aN}$ ($\times 10^{-9}$)	m_a (meV)	f_a ($\times 10^8$ GeV)
ρ_0	only NN	0.81	21.02	2.71
	$\pi N + NN$	0.46	11.99	4.75
$\rho_0/2$	only NN	0.93	24.11	2.36
	$\pi N + NN$	0.42	10.96	5.20

Bound on the effective axion-nucleon coupling $\bar{g}_{aN}$ for KSVZ axions.


Axion-Like Particles from Supernovae: the photon coupling

ALP production channels


G. Lucente, PC *et al.*, JCAP **12** (2020), 008

ALPs are coupled with photons and are produced by:

Primakoff conversion




Inverse Decay




SN1987A ALP bound

Nice complementarity with other bounds



Can ALP revitalize the SN shock?

Massive ALP could decay inside the SN revitalizing the shock




Energy deposited at $t_{\text{pb}} = 0.3 \text{ s}$, the red line indicates where the ALP deposit the same energy as neutrinos


Direct signatures from the
Diffuse SN ALP Background

SN axion phenomenology: conversion of light axions

Production




Signature



DSNALPB

F. Calore, PC *et al.*, Phys. Rev. D **102** (2020) no.12, 123005

The nucleon coupling is less constrained, larger flux with NN




DSNALPB with $g_{ap} = 1.2 \times 10^{-9}$ and $g_{a\gamma} = 5.3 \times 10^{-12} \text{GeV}^{-1}$

ALP conversion into photons

D. Horns *et al.*, Phys. Rev. D **86** (2012), 075024


The Galactic magnetic field will convert into photons both the DSNALPB and the point-like ALP flux from SN1987A (white dot)




Conversion probability for $m_a \ll E = 50 \text{ MeV}$, $g_{a\gamma} = 3 \times 10^{-13} \text{ GeV}^{-1}$

Fermi-LAT data

Skymap of gamma-rays observed by Fermi-LAT




The ALP signal



The bound


F. Calore, PC *et al.*, Phys. Rev. D **105** (2022) no.6, 063028.

The bound is stronger than CAST and can be improved by future γ -ray measurements




SN axion phenomenology: decay into electron-positron pairs

Production



Signature




$a \rightarrow e^+ e^-$ is not invisible




Positrons lose energy in $10^3 - 10^6$ yrs

Electron Positron Annihilation


- Direct annihilation




- Annihilation via positronium (Ps) formation



Annihilation line
 $E = 511$ keV




Positronium continuum
 $E < 511$ keV



Is it possible to explain **a fraction** of the 511 keV line with ALPs?


Agaronyan, F. A., and A. M. Atayan, 1981, Sov. Astr. Letters 7, 395

Let's compare with SPI data...



Very good agreement for the vertical distribution...

... much less agreement with the horizontal one




ALPs escaping from the Galaxy

Positrons trapped in the intergalactic medium ($B \sim \text{nG}$) annihilate in $\sim \text{Gyr}$ and photons are redshifted



Extragalactic X-ray diffuse flux


The extragalactic flux is redshifted, no more 511 keV line




Diffuse flux for $g_{ae} = 7 \times 10^{-21}$

Overview plot

F. Calore, PC *et al.*, Phys. Rev. D **104** (2021) no.4, 043016



Conclusions



"Always the last place you look!"