

1222·2022
800
ANNI



UNIVERSITÀ
DEGLI STUDI
DI PADOVA

arQus
European University Alliance

Ivana Batkovic

AXION-LIKE PARTICLES SEARCHES

Gamma arQus school, UIB, September 2022

SUMMARY

1. What are axion-like particles (**ALPs**)?
2. Motivation – interaction with photons
3. Experimental search for axion-like particles (**ALPs**)
4. Imaging atmospheric Cherenkov telescopes – **MAGIC**
5. **HANDS - ON**
 - ALPs study – **NGC1275**
 - a. Data selection
 - b. Spectrum analysis – fitting
 - c. ALPs analysis – photon survival probability
 - d. **Likelihood** calculations – ALPs constraints



WHAT ARE AXION-LIKE PARTICLES (ALPS)?

- Pseudoparticles of spin 0 – pseudobosons
- Solution to the strong CP problem in Standard model of particles - consequence of breaking of pseudosymmetry – **AXION**
- Generalisation of axion → axion – like particles
- Still not found! Would be a great success (free visit to Stockholm)
- In special cases they are great candidates for Dark Matter!





MOTIVATION-INTERACTION WITH PHOTONS

- Photon-ALP mixing in **the external magnetic field**
 - experimental search
 - Explanation of the irregularities in the AGN spectra?
 - Increasing the transparency of the Universe to the VHE gamma rays?
- Experimental search: helioscopes, haloscopes, **IACTs...**





Figure 1: Feynman diagram of photon-ALP interaction



MOTIVATION-INTERACTION WITH PHOTONS




Figure 2: Constraints in the ALPs parameter space, credit: <https://cajohare.github.io/AxionLimits/docs/ap.html>



MOTIVATION-INTERACTION WITH PHOTONS

- **Observable effect 1:**
 - **Irregularities (wiggles)** in the spectrum of astrophysical objects – un-explained so far – failed attempts to explain them with the effects of EBL (Extragalactic background light – all the radiation accumulated throughout the history of the Universe) – possible solution:

ALPs!





Figure 3: Spectrum fit of PKS2155-304 w & w/o ALPs, [arXiv:1311.3148](https://arxiv.org/abs/1311.3148)




MOTIVATION-INTERACTION WITH PHOTONS

- **Observable effect 2:**
 - **Spectral hardening:** increase of the photon count on the TeV energies - increasing the transparency of the Universe to the VHE gamma rays ($E > 200$ GeV) - EBL models not sufficient to explain it
 - possible solution: **ALPs!**



MOTIVATION-INTERACTION WITH PHOTONS

1. Mixing in the **blazar + jet**



4. Mixing in the **Milky Way**

2. Mixing in the **galaxy cluster**

3. Mixing in the **extragalactic space +**
($\gamma + \gamma \rightarrow e^+ + e^-$)

Figure 6: Photon-ALP mixing in the magnetic field, credit: [arXiv:0905.3270](https://arxiv.org/abs/0905.3270)

KNOWLEDGE OF THE **MAGNETIC FIELDS** IS FUNDAMENTAL FOR PRODUCING THE **ALPS** MODELS!

MOTIVATION-INTERACTION WITH PHOTONS

- We are searching for irregularities that occur around the critical energy:

$$E_{crit} = 2.5 \text{ GeV} \frac{|m_{a,neV}^2 - \omega_{pl,neV}^2|}{G_{11} B_{\mu G}}$$

- Using the **GammaALPs** code by M. Meyer: <https://gammaalps.readthedocs.io>
 - Solves the equations of motion of photon-**ALP** system
 - Inputs: **magnetic field** models, EBL model, mass of **ALPs**, coupling to photons
- Parameter space to be searched ($m_a, g_{a\gamma}$) is determined by telescope's energy range, for **IACTs**;

$$10^{-9} \text{ neV} \leq m_a \leq 10^{-6} \text{ neV}$$

$$10^{-12} \text{ GeV}^{-1} \leq g_{a\gamma} \leq 5 \times 10^{-10} \text{ GeV}^{-1}$$

MOTIVATION-INTERACTION WITH PHOTONS

- **Photon survival probability:** probability that once emitted photon will be detected with the instrument
- Needed to create models of photon-**ALP** interaction effect on the photon flux from a gamma-ray source

HANDS-ON SPOILER




Figure 5: Photon survival probability,
(<https://github.com/me-manu/gammaALPs>)

EXPERIMENTAL SEARCH FOR ALPS




Figure 6: Constraints in the ALPs parameter space, astro closeup, credit: <https://cajohare.github.io/AxionLimits/docs/ap.html>

- **Helioscopes:** looking for solar axions, in e.g. CAST
- **Haloscopes:** resonating microwave cavities looking for dark matter axions, in e.g. ADMX
- **Astrophysical constraints:** X-rays, γ -rays...(supernova, galaxy clusters, blazars...)

EXPERIMENTAL SEARCH FOR ALPS

• Helioscopes




Figure 7: CAST experiment scheme, Nat. Phys., 13, 584–590 (2017)

Most stringent
constraint on the axion
coupling up to date
(Nature Physics, 13, 584–
590 (2017)):
 $g_{a\gamma} < 6.6 \times 10^{-11} \text{GeV}^{-1}$

- Most famous example: **CAST**, future: IAXO
- Uses a 9.26 m long superconductive magnet capable of producing magnetic field up to 9.5 T (former LHC magnet) to detect photons produced in Primakoff conversion of axions in the Sun's core

EXPERIMENTAL SEARCH FOR ALPS

- **Haloscopes**

- Most famous examples: The Axion Dark Matter eXperiment (**ADMX**), MADMAX, HAYSTAC
- Uses microwave cavity inside of a large superconducting magnet (8 T) with the possibility of shifting the frequency to search for axions with masses around μeV

Ruled out dark matter axions
(Phys.Rev.Lett., 127, 26, 261803, 2021):
 $3.3 \mu\text{eV} < m_\alpha < 4.2 \mu\text{eV}$




Figure 8: ADMX experiment scheme, Nature 590, 226-227 (2021)

EXPERIMENTAL SEARCH FOR ALPS

- **Astrophysical constraints**

- In example: IACTs (Imaging Atmospheric Cherenkov Telescope): Veritas, HESS, MAGIC

- Constraints so far:

H.E.S.S.: Abramowski, A. et al.: *Phys. Rev. D* 2013, *88*, 102003.

FERMi-LAT: Ajello, M. et al.: *Phys. Rev. Lett.* 2016, *116*, 161101.

CTA: Abdalla, H. et al.: *J. Cosmol. Astropart. Phys.* 2021, *2021*, 48




Figure 9: LST1 telescope

IMAGING ATMOSPHERIC CHERENKOV TELESCOPES (IACTS)

- **Observing extensive atmospheric showers**

- The Cascades of subatomic particles in the atmosphere → Cherenkov light
- Detecting gamma-rays in the energy range of 25 GeV - 100 TeV
- Field of view $\sim 3.5^\circ$
- Angular resolution $\sim 0.1^\circ$ (energy-dependent)




Figure 10: Detection of atmospheric showers, CTA Observatory

IACTS – MAGIC EXPERIMENT

- Two Imaging Atmospheric Cherenkov telescopes located in Observatory Roque del Muchachos on the Canary island of La Palma
- 17 m diameter
- Operating since 2003, in stereo mode from 2009
- At the altitude of ~ 2240 m
- International collaboration of about 300 members from 13 countries




Figure 11: MAGIC telescopes,
Credit: Chiara Righi

ALPS ANALYSIS OF IACT DATA - KEYWORDS

- **Active galactic nuclei** – active (accreting) supermassive black hole in the center of a galaxy
- **γ -ray flux** – rate of γ -rays per unit area ($cm^{-2}s^{-1}$)
- **Gammapy** – open-source Python package for gamma-ray astronomy
- **Smooth function – PWL** (power-law), **EPWL** (power law with exponential cutoff), LP (log parabola), **ELP** (log parabola with exponential cutoff)

ALPS ANALYSIS OF IACT DATA – STEPS

1. Choose the source of the gamma-rays
2. Use the **gammapy** to obtain the source spectrum
3. Fit it with a smooth function (**PWL, EPWL, LP, ELP...**)
4. Calculate the photon survival probability using the **GammaALPs** package
5. Multiply the “**intrinsic fit**” with the photon survival probability
6. Evaluate the fit using the binned likelihood function and search for the nuisance parameters that maximise the likelihood
7. Set the constraints on the **ALPs** parameter space

CHOICE OF THE SOURCE

- a. Extragalactic source of radiation – **ACTIVE GALACTIC NUCLEI**
- b. Gamma-ray radiation – **Cherenkov telescopes**
- c. Larger data base → more precise analysis and radiation spectrum
- d. High flux states of the source: **flaring states**

Our choice:

NGC1275 in the center of the Perseus galaxy cluster
(distance: $z=0.01 \rightarrow 71,774,000$ pc)




Figure 12: Perseus galaxy cluster, credit: NASA/CXC/SAO/E.Bulbul, et al.

IACT SPECTRAL ANALYSIS AND FITTING THE SPECTRUM TO A SMOOTH FUNCTION

- **Gammapy** package

- γ -ray flux: $\Phi = \frac{d^2 N_\gamma}{dS dt}$

- Differential energy spectrum: $\frac{d\Phi}{dE} = \frac{d^3 N_\gamma}{dS dt dE}$

- Spectral energy distribution (SED): $E^2 \frac{d\Phi}{dE}$

- Fitting the SED to an EPWL

- HANDS-ON




Figure 13: SED with intrinsic fit (EPWL)

PHOTON SURVIVAL PROBABILITY

- Using the **GammaALPs** code – HANDS-ON
- Photon survival probability:

$$P_{\gamma\gamma} = \sin^2(2\theta) \sin^2 \left[\frac{g_{\alpha\gamma} B s}{2} \sqrt{1 + \left(\frac{\varepsilon}{E_\gamma} \right)^2} \right]$$

- Absorbed (with ALPs) spectrum:

$$EPWL \times p_{\gamma\gamma}$$




Figure 14: Comparison of "intrinsic" and "ALP absorbed" spectrum

BINNED LIKELIHOOD

- Binned likelihood:

$$\mathcal{L}(\theta, b) = \mathcal{L}(m_a, g_{a\gamma}; B, \Gamma, \Phi_0, E_c | b)$$

- We search for the nuisance parameters (Γ, Φ_0, E_c) that maximise the likelihood
- In the case of **ALPs**, due to the unknown **magnetic field**, random magnetic field realisations have to be employed to calibrate the test statistics for excluding the **ALPs** parameters.

CONSTRAINTS ON THE ALPS PARAMETER SPACE

- STANDARD PROCEDURE:

Use the Wilks' Theorem to convert a TS value into a significance with which one can reject the null hypothesis

CONSTRAINTS ON THE ALPS PARAMETER SPACE

- STANDARD PROCEDURE:

Use the Wilks' Theorem to convert a TS value into a significance with which one can reject the null hypothesis

- Not applicable in this situation!

CONSTRAINTS ON THE ALPS PARAMETER SPACE

- STANDARD PROCEDURE:

Use the Wilks' Theorem to convert a TS value into a significance with which one can reject the null hypothesis

- Not applicable in this situation!

Reasons:

1. The spectral irregularities do not scale linearly with the ALP parameters
2. Under the null hypothesis, the likelihood values are independent of the magnetic-field realizations.
3. Photon-ALP oscillations are completely degenerate in coupling and magnetic field

CONSTRAINTS ON THE ALPS PARAMETER SPACE

SOLUTION:

SIMULATIONS OF THE DATASETS

- How is the TS distributed in our case, how in the case of the null hypothesis and how if we include the effects of the ALPS?
- HANDS ON – SNEAK PEEK





Figure 15: Distribution of the test statistics in the case of null hypothesis

CONSTRAINTS ON THE ALPS PARAMETER SPACE



WHAT ABOUT
AFTER THE
CALIBRATION??

Figure 16: Constraints on the ALPs parameter space **BEFORE** the calibration of the TS

FUTURE PROSPECTS ON ALPS SEARCHES IN VHE RANGE

- CTA – projected limits
- Studies of blazars – detailed models for the jet magnetic field needed
- Improved understanding of the systematical uncertainties and statistical analysis
- ...




Figure 17: Projected limits from the CTA simulations, Abdalla, H. et al., doi:10.1088/1475-7516/2021/02/048

THANK YOU FOR THE
ATTENTION!

IT'S HANDS ON TIME!

HANDS ON

GammaALPs notebooks:

`/arqus_school_2022/alps/gamma_intro/alps/Py_models/gammaALPs_NGC1275.ipynb`

Gammapy & **ALPs** analysis:

`/arqus_school_2022/alps/gamma_intro/alps/ngc1275_alps_analysis.ipynb`